

QCD PHYSICS AT LEP

I.M. Dremin

Lebedev Physical Institute, Moscow

1. the energy dependence of mean multiplicities,
2. oscillations of cumulant moments of multiplicity distributions as functions of their rank,
3. difference between quark and gluon jets,
4. the hump-backed plateau of inclusive rapidity distribution and energy dependence of its maxima,
5. difference between heavy- and light-quark jets,
6. color coherence in 3-jet events,
7. intermittency and fractality
8. the energy behavior of higher moments of multiplicity distributions,
9. subjet multiplicities,
10. jet universality.

Recent review papers:

Dremin, Gary, Phys. Rep. 349 (2001) 301

Dremin, Physics-Uspekhi 172 (2002) 551
(www.ufn.ru → May 2002 issue)

Theory

The generating functional:

$$G(\{u\}, \alpha_0) = \sum_n \int d^3 k_1 \dots d^3 k_n u(k_1) \dots u(k_n) P_n(k_1, \dots, k_n; \alpha_0)$$

$P_n(k_1, \dots, k_n; \alpha_0)$ - the probability density of exclusive production of particles with momenta k_1, \dots, k_n at the initial energy α_0 .

$u(k)$ - an auxiliary function

For $u(k) = u = \text{const}$, $P_n(\alpha_0)$ - the multiplicity distribution

$$G(\alpha_0) = \sum_n u^n P_n(\alpha_0) \quad \text{- the generating function}$$

Symbolical equation (gluons) \rightarrow system of two eqs for q and g jets

$$G' \sim \int d_s K [G \otimes G - G] ds$$

phase space

Cascade evolution (G') is determined by 3-gluon vertex (\otimes) (production) and escape of a single gluon (G) weighted by coupling strength α_s and a splitting function K defined by the interaction Lagrangian

$$G'(y) = \int_0^1 dx K(x) y^2 [G(y + \ln x) G(y + \ln(1-x)) - G(y)]$$

$$\underline{y}^2 = \frac{2 N_c \alpha_s}{\pi} ; \quad \underline{y} = \ln \frac{P \theta}{Q_0} ; \quad P - \text{jet momentum, } \theta - \text{opening angle}$$

$$K = \frac{1}{x} - (1-x)[2-x(1-x)]$$

LO \equiv DLA $\rightarrow [\alpha_s \ln^2 s]^n$ terms summed

NLO \equiv MLLA $\rightarrow [\alpha_s \ln s]^n$

2NLO, 3NLO ... \rightarrow kinetic equation (?)

Gluodynamics

LO (DLA) $\sum_n [\alpha_s \ln^2 s]^n$

$K(x) \approx \frac{1}{x}$; $\alpha_s \approx \text{const. (fixed)}$ or $\alpha_s \sim \frac{1}{y} \text{ (running)}$

$$G(y + \ln(1-x)) \approx G(y)$$

$$[\ln G(y)]' \approx \gamma_0^2 \left\{ \frac{dx}{x} [G(y + \ln x) - 1] \right\}$$

$$[\ln G(y)]'' \approx \gamma_0^2 [G(y) - 1] \quad - \text{LO equation}$$

b.c.: $G(0, u) = u$; $G(y, 1) = 1 (= \sum_n P_n)$

NLO (MLLA) $\sum_n [\alpha_s \ln s]^n$

Renormgroup approach - A. Mueller (1984)

No systematic perturbative expansion
till 1993 → Dremin (1993)

↓
"Exponential" perturbative series -
- modified perturbative expansion (MPE)

Taylor series expansion \equiv MPE

Exact solution for fixed coupling - Dremin, Hwa
Scaling → power behavior (1994)

Parton and dipole approaches in QCD

I.M. Dremin¹, P. Edén²

¹Lebedev Physical Institute, Moscow 119991, Russia

²Nordita, Copenhagen, DK-2100, Denmark

Factorial moments

$$F_q = \frac{d^q G(u)}{du^q} \Big|_{u=1}; \quad \text{Cumulant moments} \quad K_q = \frac{d^q \ln G(u)}{du^q} \Big|_{u=1}; \quad H_q = \frac{K_q}{F_q} \quad (1)$$

$$G(u, y) = \sum_n u^n P_n(y), \quad G' \sim \int \alpha_S K[G \otimes G - G] d\Omega. \quad (2)$$

init. cond. $P_n = \delta_{n1}$; $G_0 = u$. (3)

$$G'_G = \int_0^1 dx K_G^G(x) \gamma_0^2 [G_G(y + \ln x) G_G(y + \ln(1-x)) - G_G(y)] \\ + n_f \int_0^1 dx K_G^F(x) \gamma_0^2 [G_F(y + \ln x) G_F(y + \ln(1-x)) - G_G(y)], \quad (4)$$

$$G'_F = \int_0^1 dx K_F^G(x) \gamma_0^2 [G_G(y + \ln x) G_F(y + \ln(1-x)) - G_F(y)], \quad (5)$$

$$y = \ln(p\Theta/Q_0) = \ln(2Q/\underline{\underline{Q_0}})$$

$$\underline{\underline{\gamma_0^2}} = \frac{2N_c \alpha_S}{\pi},$$

$$\alpha_S(y) = \frac{2\pi}{\beta_0 y} \left(1 - \frac{\beta_1}{\beta_0^2} \cdot \frac{\ln 2y}{y} \right) + O(y^{-3}), \quad (7)$$

$$K_G^G(x) = \frac{1}{x} - (1-x)[2-x(1-x)], \quad (8)$$

$$K_G^F(x) = \frac{1}{4N_c} [x^2 + (1-x)^2], \quad (9)$$

$$K_F^G(x) = \frac{C_F}{N_c} \left[\frac{1}{x} - 1 + \frac{x}{2} \right], \quad (10)$$

Energy conservation

Cut-off parameter (non-perturb.)
[LPHD]

DG (01)

+ LPHD (1)

Experiment & QCD

1. $\langle n(E) \rangle$

- mean multiplicity

$$\left. \frac{dG}{du} \right|_{u=1} = \sum n P_n = \langle n(E) \rangle \sim e^{\int_0^y \gamma(y') dy'}$$

Pert. expansion of exponent: MPE $\rightarrow \gamma = \gamma_0(1 - a_1 \gamma_0 - a_2 \gamma_0^2 - \dots)$

Prediction: $\langle n \rangle \sim \exp[c \sqrt{\ln s}]$

DLA, running coupling $[\gamma_0 \sim \frac{1}{\sqrt{s}}]$
MLLA \rightarrow factor $\ln^{-d} s \dots$

Hydrodynamics; fixed coupling QCD \rightarrow power behavior

Multiperipheral; flat rapidity plateau \rightarrow log behavior

Fig. 1

(Feynman)

2. H_q -moments

- shape of the distribution

$$\left. \frac{1}{\langle n \rangle^q} \frac{d^q G}{du^q} \right|_{u=1} = \frac{\langle n(n-1) \dots (n-q+1) \rangle}{\langle n \rangle^q} = F_q \quad \text{factorial moments}$$

$$\left. \frac{1}{\langle n \rangle^q} \frac{d^q \ln G}{du^q} \right|_{u=1} = K_q \quad \begin{matrix} \text{cumulant} \\ \text{moments} \end{matrix} \quad \begin{matrix} \text{"genuine"} \\ \text{(correlations)} \end{matrix}$$

$$H_q = \frac{K_q}{F_q}$$

Prediction: DLA $\rightarrow H_q = 1/q^2$

New effect: MLLA \rightarrow minimum at $q \approx \frac{24}{11 \gamma_0} + 0.5 \approx 5$
predicted 2NLO... \rightarrow oscillations

Poisson: $H_q \equiv 0$


Negative Binomial: $H_q = \frac{\Gamma(q) \Gamma(k+1)}{\Gamma(k+q)} = k B(q, k) > 0$

Fig. 2

(clusters) attraction \leftrightarrow repulsion (separated jets)
 (nothing in NBD!)

Dremin
1993

Dremin
Nechitai
1993



- Mueller
- Webber
- Dremin, Nechitailo
- Capella, Dremin, Gary, Nechitailo, Tran Thanh Van

Perturbative solutions

→ Taylor series expansion.

$$r = r_0(1 - r_1 y_0 - r_2 y_0^2 - r_3 y_0^3) + O(y_0^4) \quad (*)$$

$$\gamma = \gamma_0 (1 - a_1 \gamma_0 - a_2 \gamma_0^2 - a_3 \gamma_0^3) + O(\gamma_0^5) \quad (**)$$

$$\mathcal{L}_S = \frac{2\pi}{\beta_0 y} \left[1 - \frac{\beta_1}{\beta_0^2} \cdot \frac{\ln 2y}{y} \right] + O(y^{-3}) \xrightarrow{\text{2-loop}} ; \quad y_0 = \sqrt{\frac{2N_{cols}}{\pi}} \quad (\text{at } y_0)$$

$$\beta_0 = \frac{11N_c - 2n_f}{3}; \quad \beta_1 = \frac{51N_c - 19n_f}{3}$$

tends to 0 at $y \rightarrow \infty$
 as $y^{-\frac{1}{2}}$ (as freedom energies)

Analytic formulas for a_i, r_i [$\rightarrow = 0$ in SUSY-QCD]

Numerical values:

$$r_0 = \frac{C_A}{C_F} = \frac{9}{4} \quad \rightarrow LO \equiv DLA \quad (NLO \equiv MLLA)$$

n_f	r_1	r_2	r_3	a_1	a_2	a_3
3	0.185	0.426	0.189	0.280	-0.379	0.209
4	0.191	0.468	0.080	0.297	-0.339	0.162
5	0.198	0.510	-0.041	0.314	-0.301	0.112
	NLO_F	$NNLO_F$	$3NLO_F$	NLO	$NNLO$	$3NLO$
	$2NLO$	$3NLO$	$4NLO$			

Figs. Ratio : $2.25 \rightarrow 2.03 \rightarrow 1.77 \rightarrow 1.74$ at Z^0 for $\gamma_0 = 0.5$, $n_f = 4$

→ Anomalous dimension γ determines energy dependence:

$$\langle n_G \rangle = A \exp \left[\int^y \gamma(y) dy \right]$$

Fig.

\downarrow y_0
non-perturbative normalization constant
(fixed parameter)


Perturbative vs non-perturbative regions in Eqns:

$\int_{\text{out}}^{\text{in}} \dots dx$ includes both P and NP-regions but admits purely perturbative solutions $(*)$ and $(**)$ } 2

$\int_{e^{-y}}^{1-e^{-y}}$ includes just P region ($Q \geq Q_0$) but gets NP-terms $\sim e^{-y} \sim \frac{1}{Q}$ (no analytic solution)

$$r_{\text{ch.}} = \frac{\langle n_{\text{ch.}} \rangle_{\text{gluon}}}{\langle n_{\text{ch.}} \rangle_{\text{quark}}} = 1.51 \pm 0.02 \pm 0.05$$

$E_{\text{jet}} \approx 41.6 \text{ GeV}$



Energy dependence:

$$\langle n_G \rangle = A y^{-\alpha c^2} \exp \left\{ 2c\sqrt{y} + \frac{c}{\sqrt{y}} \left[2\alpha_2 c^2 + \frac{\beta_1}{\beta_0^2} (\ln 2y + 2) \right] + \frac{c^2}{y} \left[\alpha_3 c^2 - \alpha_1 \frac{\beta_1}{\beta_0^2} (\ln 2y + 1) \right] \right\}$$

$$c = \sqrt{\frac{4N_c}{\beta_0}} \quad ; \quad y \rightarrow \ln \frac{E_{cm}}{\Lambda}$$

Two parameters: A, Λ

($c=1.2$ for $n_g=4$)

$$\delta(y) = \frac{c}{\sqrt{y}} \left[2\alpha_2 c^2 + \frac{\beta_1}{\beta_0^2} (\ln 2y + 2) \right] + \frac{c^2}{y} \left[\alpha_3 c^2 - \alpha_1 \frac{\beta_1}{\beta_0^2} (\ln 2y + 1) \right] \approx \text{const}$$

(at present energies!)

↓
NNLO+3NLO

Conclusion: NNLO+3NLO renormalize A , and the energy dependence is well approximated by NLO \equiv MLLA

$$\langle n_F \rangle = \frac{\langle n_G \rangle}{r(y)} = \frac{\langle n_G \rangle}{r_0(1 - r_1 \gamma_0 - r_2 \gamma_0^2 - r_3 \gamma_0^3)}$$

rewrite it in the exponent
↓

$$\langle n_F \rangle = \frac{A}{r_0} y^{-\alpha c^2} \exp \left\{ 2c\sqrt{y} + \frac{c}{\sqrt{y}} \left[r_1 + 2\alpha_2 c^2 + \frac{\beta_1}{\beta_0^2} (\ln 2y + 2) \right] + \frac{c^2}{y} \left[\alpha_3 + r_2 + \frac{r_1^2}{\alpha} - \alpha_1 \frac{\beta_1}{\beta_0^2} (\ln 2y + 1) \right] \right\}$$

- 1) No term with r_3 ! It would need α_3 to be self-consistent in pQCD
Conventional definition of LO, NLO, ...
- 2) Large value of $2r_2/r_1 \sim 5$ (!) (compare with $\alpha_3 \sim 0.15 \div 0.2$)
- 3) The same dependence of $\langle n_G \rangle$ and $\langle n_F \rangle$ in MLLA \equiv NLO!

$$\gamma_F = \gamma - \frac{r'}{r} \quad ; \quad r' \sim \gamma_0' \sim \gamma_0^3 \rightarrow \text{SLOPES} !$$

$$\gamma_F < \gamma$$

↑
NNLO correction in γ_F

$$\gamma_{F_{NLO}} = \gamma_{NLO}$$

$$r < r^{(1)} < r^{(2)} < \frac{C_A}{C_F} = 2.25$$

$$r^{(1)} = \frac{\langle n_G \rangle'}{\langle n_F \rangle}, \quad \begin{matrix} \text{ratio} \\ \text{of} \\ \text{slopes} \end{matrix}$$

$$r^{(2)} = \frac{\langle n_G \rangle''}{\langle n_F \rangle''} \dots \begin{matrix} \text{curvat} \end{matrix}$$

→ Figs.

NP effects






Figure 2: The average multiplicities of gluon (G) and quark (F) jets in the NLO and 3NLO approximations, compared to data [13]. For the theory, the gluon jet normalization is fit to the data; the normalization of the gluon jet curve fixes the normalization of the quark jet curve. The theoretical results are obtained using $n_f = 4$.

e+e- 91./91.5 GeV DELPHI Coll.



q Dr. Arena et al Phys. Lett.

$$q_{\min} \approx \frac{1}{h_i \gamma_0} + \frac{1}{2} \approx 5; \quad h_i = \frac{11}{24}; \quad \gamma_0 \approx 0.48 \text{ at } Z^0 \quad B336(1994) 119 \\ + [SLD \text{ Collaboration}]$$



12-95

8058A4

- Dremin (93)
- Dremin, Nechitailo (93)
- Lupia (98)

3. q & g jets

$r = \frac{\langle n_g \rangle}{\langle n_q \rangle}$ and $\gamma_G, \gamma_F \rightarrow$ different mean multiplicities and anomalous dimension

Prediction:

$$\text{DLA} \rightarrow r_0 = \frac{C_A}{C_F} = \frac{9}{4}; \quad r = \frac{9}{4}(1 - r_0 \gamma_0 - r_0^2 \gamma_0^2 - \dots)$$

Fig. 3

$$\gamma_G \approx \gamma_F \quad \text{MLLA} \rightarrow \gamma_G = \gamma_F \quad \gamma_F = \gamma_G - \frac{r}{r_0} < \gamma_G$$

Fig. 3a \rightarrow **Fig. 1**

4. Hump-backed plateau (vs Feynman flat plateau)

Inclusive distribution \rightarrow eq. for generating functional

Prediction:

- 1) Maxima ($\xi = \ln \frac{1}{x} = \ln \frac{E_j}{P}$) \rightarrow **Fig. 4** [angular ordering and color coherence]
- 2) Shape \approx Gaussian \rightarrow **Fig. 5** [MLLA]
- 3) Maxima positions and energy dependence \rightarrow **Fig. 6** (widths...) **Fig. 6a**


5. Heavy- and light-quark jets

Analogy: Bremsstrahlung of e and μ ;
role of mass in the propagator $\frac{1}{k^2 + m^2}$

Prediction:


Energy-independent difference of companion mean multiplicities of heavy- and light-quark jets of equal energy

Fig. 7 \rightarrow compare to "naive" model



- Azimov, Dokshitzer, Khoze (82)
- Dokshitzer, Fadin, Khoze (82, 83)
- Bassetto, Ciafaloni, Marchesini, Mueller (82)

Peak Position and Width of ξ of Charged Particle Spectra



- MLLA+LPHD fit describes data well
- No deviation from e^+e^- behaviour
 - Fong, Webber (89-91)
 - Dokshitzer, Khoze, Troyan (91)




Figure 10: Peak position ξ^* of the inclusive ξ distribution plotted against di-jet mass $\times \sin(\Theta)$ in comparison with the MLLA prediction (central curve); also shown are the double logarithmic approximation (lower curve with asymptotic slope $\xi^* \sim Y/2$) and expectation from cascade without coherence. Result by CDF Collaboration.⁹²

data at full angle $\Theta = \pi/2$ from e^+e^- and ep collisions whereby the variable $Y = \ln(P \sin \Theta / Q_0)$ has been used. The data scatter around the expected curve (79) for $n_f = 3$. Taking instead the scaling variable $Y = \ln(2P \sin(\Theta/2) / Q_0)$ the full angle data would be shifted to the right by a factor $2 \sin(\pi/4) \sim 1.4$. This would correspond essentially to a change of the next-to-next-to-leading order term in (79) but would not change the slope.


The slope is nicely confirmed and the leading DLA contribution ($\xi^* \sim Y/2$) is shown for comparison as the lower curve in Fig. 10, adjusted in height; the upper curve represents the spectrum for the incoherent cascade which peaks near the maximum ($\xi^* \sim Y$). Apparently the data support the prediction from the parton cascade with suppression of soft particles due to coherent gluon emission in a large energy range $2E_{jet} \sin \Theta \sim 4 - 300$ GeV.

The analytical results for the particle spectrum near the soft limit (90) are nicely confirmed by the data. In these calculations the model (35) for mass effects has been used. The experimental data from the available range

Specifics of heavy-quark jets

- Nothing very special in $H_q \rightarrow$ similar oscillations
(Dr., Nechitailo)
- Accompanying gluon radiation

$$\frac{d\sigma_b/d\theta}{d\sigma_{uds}/d\theta} \approx \frac{1}{3} \frac{\theta^4}{(\theta^2 + \frac{m_b^2}{E^2})^2}$$




Dr. JETP Lett. 30, 140 (1979) → Chir. gluons;


Ring-like (Dr. JETP Lett. 34 (1981) 534) ... t-quark

Dead-cone (Schechter, Dokshitzer, Khoze, Kotke
Phys. Rev. Lett. 69 (92) 3029)

$\langle n_a \rangle_b < \langle n_a \rangle_{uds}$; does not depend on W .


DELPHI data (unpublished)





- Azimov, Dokshitzer, Khoze (82) E_{cm} (GeV)
- Dokshitzer, Fadin, Khoze (82, 83)
- Petrov, Kisselov (88, 95)
- Dremin (79, 81)

6. Color coherence



Prediction:

1. Particle flows \rightarrow enlarged in the partons directions, depleted in-between

2. $q\bar{q}$ valley/ $q\bar{q}$ valley = 2.4 (theor) 2.23 ± 0.37 (exper.)

Fig. 8

$$3. R_\gamma = \frac{N_{q\bar{q}}(q\bar{q}g)}{N_{q\bar{q}}(q\bar{q}\gamma)} = 0.61 \text{ (theor)} \quad 0.58 \pm 0.06 \text{ (exper.)}$$

$$4. \frac{\log g}{\log \theta} \quad \theta \rightarrow 0 \quad n_{q\bar{q}g} \downarrow, n_\perp \downarrow$$

5. azimuthal correlations

7. Intermittency and fractality

Self-similarity of the parton cascade

Analogy with turbulence \rightarrow „jets inside jets inside.”

QCD Prediction:

Moments increase for bins decreasing:

a) linearly on log-log plot for large bins,


b) flattening at small bins due to α_s running.

a) \rightarrow monofractal ; b) \rightarrow multifractal

Slopes at different ranks $q \rightarrow$ Renyi dimensions



Fig. 9

Qualitatively OK, quantitatively - not
but DLA + some MLLA terms



- Andersson, Gustafson, Sjöstrand (80)
- Azimov, Dokshitzer, Khoze, Troyan (85,86)

"Genuine" multiplicity correlations in
ring regions



- (a) Dokshitzer, Dremin (93)
- (b) Brak, Peschanski (94)
- (c) Ochs, Wosiek (92, 93)

disagreement for $\Lambda = 0.16$

- Bialas, Peschanski (86)
- Dremin (87)

$\Lambda = 0.16 \text{ GeV}$

$(n_f = 3)$

8. Higher moments

Prediction:

1. Increase with rank and energy ~OK
2. Asymptotic values

$$\text{DLA} \rightarrow F_2^G = \frac{4}{3}; F_2^F = \frac{7}{4} \quad -\text{very far from } Z^\circ \text{-values}$$

3. At Z° :

$\text{MLLA} \rightarrow F_2^G \approx 1.039$	1.023 (exper.)
$\rightarrow F_2^F \approx 1.068$	1.082 (exper.)

Failure of analytic approach at higher orders!

Soft partons play a crucial role in correlations.
(Dremin, Lam, Nechitailo (2000)) ($p_T \rightarrow$ Dremin, Eden (2001))

9. Subjet multiplicities

Increase the resolution and get more subjets.

For very high resolution \rightarrow final hadrons.

Prediction:

- 1) Asymptotic ratio (DLA) of subjet multiplicities in 3- and 2-jet events:

$$\frac{n_{3j}^{sj}}{n_{2j}^{sj}} = \frac{2C_F + C_A}{2C_F} = \frac{17}{8} \longleftrightarrow < 1.5 \text{ (exper. at } Z^\circ)$$

Depletion due to color coherence.

- 2) Subjet multiplicities in separated q and q jets



Fig. 10

10. Jet universality

Prediction:

Universal in different processes.

See Figs. above and next talks



- Dokshitzer, Khoze, Troyan (83)
- Ochs, Wosiek (93)
- Seymour (96)

Conclusions

**Success
of
Analytic
pQCD**

1. Qualitative predictions - OK!
2. Quantitatively $\rightarrow 20\div 10\%$ accuracy

Computer solution & Monte Carlo models lead to better fits

Main problems:

- Soft partons and non-perturbative terms (near phase space boundary)
 - Asymptotical nature of perturbative expansion, convergence and large value of the expansion parameter ($\gamma_0 \sim 0.5$)
 - Probabilistic scheme limitations
-

event shapes

OUTLOOK

RHIC, LHC, ... - multiparticle production
 $\langle \frac{dN}{dy} \rangle \gtrsim 4000$ at 130 GeV Au-Au

QCD predictions - closer to asymptotics -
 - better estimation of background for new physics

Event-by-event analysis - pattern recognition.
 - wavelets as an analyzing tool

Color-suppressed effects, minijet properties,
 collective flows, ring-like events, event shapes,
 vacuum (?), ...