

# What are the effects of the initial-condition fluctuations in hydrodynamical description of heavy-ion collisions?

*C.E. Aguiar<sup>1</sup>, F. Grassi<sup>2</sup>, T. Kodama<sup>1</sup>,  
T. Osada<sup>3</sup> L.L.S. Portugal<sup>1</sup>,  
O. Socolowski Jr.<sup>2</sup>, B.M. Tavares<sup>1</sup> and  
Y. Hama<sup>2</sup>*

*<sup>1</sup> Instituto de Física - UFRJ*

*<sup>2</sup> Instituto de Física - USP*

*<sup>3</sup> Ochanomizu University, Tokyo, Japan*

## Plan of presentation:

1. Purpose of the study
2. Method
3. Results
4. Conclusions

# Purpose of the study

Usual hydrodynamic approach



Symmetric and smooth initial conditions:  
(mean distributions of velocity, temperature,  
energy density, etc. )


However, our systems are not large enough.



Large fluctuations are expected.

What are the effects of the event-by-event  
fluctuation of the initial conditions?

- Are they sizable?
- Do they depend on the equation of state?
- Which are the most sensitive variables?
- . . .



Initial energy density distribution on  $z = 0$  plane of a typical Au+Au event at  $\sqrt{s} = 200 A \text{ GeV}$ , impact parameter  $b = 0$ , produced by NeXus event generator.



In **previous reports** [1], by neglecting baryon-number and strangeness conservation and using simple parametrizations for EoS [4], we showed that

- indeed they are sizeable
- and
- they do depend on the EoS.

Here, we shall present comparison of results obtained

with **fluctuating initial conditions**  
and those

with **averaged initial conditions.**

The **EoS** used is more realistic one [4], with baryon-number conservation taken into account and with an explicit inclusion of many resonances.

## Method of study

Generation of  
Initial Conditions  
**NeXus**

Equation of state

- Hadron gas
- QGP + hadron gas



Resolution of hydrodynamic  
equations  
**SPheRIO**




Computation of  
observables

$$\langle v_2 \rangle, \frac{dN}{dy}, \frac{d\sigma}{dm_T},$$

...

## Equation of State

We consider an EoS with first-order transition between QGP and resonance-gas (RG) phases, with baryon-number conservation and excluded volume effects taken into account. The strangeness conservation has not been introduced yet.



# Smoothed Particle Hydrodynamics

The main **characteristics** are

- To attach conserved quantities (baryon number, strangeness, entropy, etc. ) to small volumes called “**particles**”;
- Physical quantities are computed by averaging over **particles**, using some interpolating kernel;
- The **particle** motions are described by using **Lagrangian coordinates**.

## **Advantages:**

- No extra grid points are needed;
- The precision is controlled by the interpolating kernel and the volumes of the **particle**.



## SPH equations of motion

In the present work, besides the energy and momentum, we have chosen the **entropy** as our conserved quantity. Then, its density (in the space-fixed frame) is parametrized as

$$s^*(\mathbf{x}, t) = \sum_i^N \nu_i W(\mathbf{x} - \mathbf{x}_i(t); h) ,$$

where

$$\left\{ \begin{array}{l} W(\mathbf{x} - \mathbf{x}_i(t); h) \text{ is the normalized } \mathbf{kernel}; \\ \mathbf{x}_i(t) \text{ is the } i\text{-th. } \mathbf{particle} \text{ position, so} \\ \quad \text{the } \mathbf{velocity} \text{ is } \mathbf{v}_i = d\mathbf{x}_i/dt ; \\ h \text{ is the smoothing } \mathbf{scale} \text{ } \mathbf{parameter}; \end{array} \right.$$

and we have


$$S = \int d^3\mathbf{x} s^*(\mathbf{x}, t) = \sum_i^N \nu_i .$$

The **equations of motion** write

$$\begin{aligned} & \frac{d}{dt} \left( \nu_i \frac{P_i + \varepsilon_i}{s_i} \gamma_i \mathbf{v}_i \right) \\ & + \sum_j \nu_j \left[ \frac{P_i}{s_i^{*2}} + \frac{P_j}{s_j^{*2}} \right] \nabla_i W(\mathbf{x}_i - \mathbf{x}_j; h) = 0 , \end{aligned}$$



### Rapidity Distributions (Pb+Pb, 17.3A GeV)



### Transverse-Mass Distributions (Pb+Pb, 17.3A GeV)










Fig. 2: Event-by-event fluctuating distributions, compared with data [10]. The average distributions with corresponding dispersions are also shown.

## Rapidity Distributions (Pb+Pb, 17.3A GeV)




### Charged Particles; Au+Au (130A GeV)









Charged Particles; Au+Au (200A GeV)  
5 - 10%




Charged Particles; Au+Au (200A GeV)  
10 - 20%




Charged Particles; Au+Au (200A GeV)  
20 - 30%




Pseudo-rapidity Distributions (Au+Au; 200A GeV)  
Charged Particles




Neg. Charged Particles; Au+Au (130A GeV)  
0 - 5%




## Pseudorapidity Distributions (Au+Au, 130A GeV)



Centrality Dependence of  $v_2$  (pions; Pb+Pb, 17.3 A GeV)



$p_T$  dependence of  $v_2$  (pions; Pb+Pb, 17.3 A GeV)



$p_T$  dependence of  $v_2$  (protons; Pb+Pb, 17.3 A GeV)










Fig. 5






## Transverse Mass Distributions (Pb+Pb, 17.3A GeV)



## Transverse Mass Distributions (Pb+Pb, 17.3A GeV)



## Rapidity Distributions (Pb+Pb, 17.3A GeV)



## Conclusions and outlook

The effects of the event-by-event fluctuation of the initial conditions in hydrodynamics are large and should be considered in data analyses .

We emphasize that such fluctuations are not only due to the uncontrollable event-dependent impact-parameter fluctuation, but they appear even if the impact-parameter could be fixed and are due to the finite size of our systems.

In the present work, many important factors have not been considered : strangeness conservation, continuous emission effects, spectators, etc.

Also, there are many other observables : spectra for different species, correlations, etc.

We are working on some of them and will continue to work on the others.



## References

1. (**Preliminary**) T. Osada, C.E. Aguiar, Y. Hama and T. Kodama, nucl-th/0102011; C.E. Aguiar, Y. Hama, T. Kodama and T. Osada, *Nucl.Phys.* **A698** (2002) 639-642.
2. (**SPH**) C.E. Aguiar, T. Kodama, T. Osada and Y. Hama, *J. Phys.* **G27** (2001) 75.
3. (**NeXus**) H.J. Drescher, M. Hladik, S.V. Ostapchenko, T. Pierog and K. Werner, *J.Phys.* **G25** (1999) L91; *Nucl.Phys.* **A661** (1999) 604; H.J. Drescher, F.M. Liu, S.V. Ostapchenko, T. Pierog and K. Werner, *Phys.Rev.* **C65** (2002) 054902.
4. (**EoS**) C.M. Hung and E.V. Shuryak, *Phys. Rev.Lett.* **75** (1995) 4003. G.D. Yen, M.I. Gorenstein, W. Greiner and S.N. Yang, *Phys.Rev.* **C56** (1997) 2210; C.M. Hung and E.V. Shuryak, *Phys.Rev.* **C57** (1998) 1891.

5. (**SPS data**) F. Siklér, NA49 Collab., *Nucl. Phys. A* **661** (1999) 45c; A.M. Poskanzer and S.A. Voloshin, NA49 Collab., *Nucl. Phys. A* **661** (1999) 341c; H. Appelhäuser *et al.*, NA49 Collab., *Phys. Rev. Lett.* **80** (1998) 4136; *Phys. Rev. Lett.* **82** (1999) 2471.
6. (**RHIC data**)  
**STAR** Collab., K.H. Ackermann *et al.*, *Phys.Rev.Lett.* **86** (2001) 402; C. Adler *et al.*, *Phys.Rev.Lett.* **87** (2001) 112303;  
**BRAHMS** Collab., I.G. Bearden *et al.*, *Phys.Lett. B* **523** (2001) 227; *Phys.Rev.Lett.* **88** (2002) 202301.