

Study of tau-pair production in photon-photon collisions at LEP and limits on the anomalous electromagnetic moments of the tau lepton

The DELPHI Collaboration

J. Abdallah²⁵, P. Abreu²², W. Adam⁵¹, P. Adzic¹¹, T. Albrecht¹⁷, T. Alderweireld², R. Alemany-Fernandez⁸, T. Allmendinger¹⁷, P. Allport²³, U. Amaldi²⁹, N. Amapane⁴⁵, S. Amato⁴⁸, E. Anashkin³⁶, A. Andreazza²⁸, S. Andringa²², N. Anjos²², P. Antilogus²⁵, W-D. Apel¹⁷, Y. Arnaud¹⁴, S. Ask²⁶, B. Asman⁴⁴, E. Augustin²⁵, A. Augustinus⁸, P. Baillon⁸, A. Ballestrero⁴⁶, P. Bambade²⁰, R. Barbier²⁷, D. Bardin¹⁶, G. Barker¹⁷, A. Baroncelli³⁹, M. Battaglia⁸, M. Baubillier²⁵, K-H. Becks⁵³, M. Begalli⁶, A. Behrmann⁵³, E. Ben-Haim²⁰, N. Benekos³², A. Benvenuti⁵, C. Berat¹⁴, M. Berggren²⁵, L. Berntzon⁴⁴, D. Bertrand², M. Besancon⁴⁰, N. Besson⁴⁰, D. Bloch⁹, M. Blom³¹, M. Bluj⁵², M. Bonesini²⁹, M. Boonekamp⁴⁰, L. Booth²³, G. Borisov²¹, O. Botner⁴⁹, B. Bouquet²⁰, V. Bowcock²³, I. Boyko¹⁶, M. Bracko⁴³, R. Brenner⁴⁹, E. Brodet³⁵, P. Bruckman¹⁸, M. Brunet⁷, L. Bugge³³, P. Buschmann⁵³, M. Calvi²⁹, T. Camporesi⁸, V. Canale³⁸, F. Carena⁸, N. Castro²², F. Cavallo⁵, M. Chapkin⁴², Ph. Charpentier⁸, P. Checchia³⁶, R. Chierici⁸, P. Chliapnikov⁴², J. Chudoba⁸, U. Chung⁸, K. Cieslik¹⁸, P. Collins⁸, R. Contri¹³, G. Cosme²⁰, F. Cossutti⁴⁷, J. Costa⁵⁰, B. Crowley¹, D. Crennell³⁷, J. Cuevas³⁴, J. D'Hondt², J. Dalmau⁴⁴, T. da Silva⁴⁸, W. Da Silva²⁵, G. Della Ricca⁴⁷, A. De Angelis⁴⁷, W. De Boer¹⁷, C. De Clercq², B. De Lotto⁴⁷, N. De Maria⁴⁵, A. De Min³⁶, L. de Paula⁴⁸, L. Di Ciaccio³⁸, A. Di Simone³⁹, K. Doroba⁵², J. Drees^{53,8}, M. Dris³², G. Eigen⁴, T. Ekelof⁴⁹, M. Ellert⁴⁹, M. Elsing⁸, C. Espirito Santo²², G. Fanourakis¹¹, D. Fassouliotis^{11,3}, M. Feindt¹⁷, J. Fernandez⁴¹, A. Ferrer⁵⁰, F. Ferro¹³, U. Flagmeyer⁵³, H. Foeth⁸, E. Fokitis³², F. Fulda-Quenzer²⁰, J. Fuster⁵⁰, M. Gandelman⁴⁸, C. Garcia⁵⁰, Ph. Gavillet⁸, E. Gazis³², R. Gokieli^{8,52}, B. Golob⁴³, G. Gomez-Ceballos⁴¹, P. Goncalves²², E. Graziani³⁹, G. Grosdidier²⁰, K. Grzelak⁵², J. Guy³⁷, C. Haag¹⁷, A. Hallgren⁴⁹, K. Hamacher⁵³, K. Hamilton³⁵, S. Haug³³, F. Hauler¹⁷, V. Hedberg²⁶, M. Hennecke¹⁷, H. Herr⁸, J. Hoffman⁵², S-O. Holmgren⁴⁴, J. Holt⁸, A. Houlden²³, K. Hultqvist⁴⁴, N. Jackson²³, G. Jarlskog²⁶, P. Jarry⁴⁰, D. Jeans³⁵, K. Johansson⁴⁴, D. Johansson⁴⁴, P. Jonsson²⁷, C. Joram⁸, L. Jungermann¹⁷, F. Kapusta²⁵, S. Katsanevas²⁷, E. Katsoufis³², G. Kernel⁴³, P. Kersevan^{8,43}, U. Kerzel¹⁷, A. Kiiskinen¹⁵, T. King²³, J. Kjaer⁸, P. Kluit³¹, P. Kokkinias¹¹, C. Kourkouvelis³, O. Kouznetsov¹⁶, Z. Krumstein¹⁶, M. Kucharczyk¹⁸, J. Lamsa¹, G. Leder⁵¹, F. Ledroit¹⁴, L. Leinonen⁴⁴, R. Leitner³⁰, J. Lemonne², V. Lepeltier²⁰, T. Lesiak¹⁸, W. Liebig⁵³, D. Liko⁵¹, A. Lipniacka⁴⁴, H. Lopes⁴⁸, M. Lopez³⁴, D. Loukas¹¹, P. Lutz⁴⁰, L. Lyons³⁵, J. MacNaughton⁵¹, A. Malek⁵³, S. Maltezos³², F. Mandl⁵¹, J. Marco⁴¹, R. Marco⁴¹, B. Marechal⁴⁸, M. Margoni³⁶, J-C. Marin⁸, C. Mariotti⁸, A. Markou¹¹, C. Martinez-Rivero⁴¹, J. Masik¹², N. Mastroiannopoulos¹¹, F. Matorras⁴¹, C. Matteuzzi²⁹, F. Mazzucato³⁶, M. Mazzucato³⁶, R. Mc Nulty²³, C. Meroni²⁸, T. Meyer¹, E. Migliore⁴⁵, W. Mitaroff⁵¹, U. Mjoernmark²⁶, T. Moa⁴⁴, M. Moch¹⁷, K. Moenig^{8,10}, R. Monge¹³, J. Montenegro³¹, D. Moraes⁴⁸, S. Moreno²², P. Morettini¹³, U. Mueller⁵³, K. Muenich⁵³, M. Mulders³¹, L. Mundim⁶, W. Murray³⁷, B. Muryn¹⁹, G. Myatt³⁵, T. Myklebust³³, M. Nassiakou¹¹, F. Navarra⁵, K. Nawrocki⁵², R. Nicolaidou⁴⁰, M. Nikolenko^{16,9}, A. Oblakowska-Mucha¹⁹, V. Obraztsov⁴², A. Olshevski¹⁶, A. Onofre²², R. Orava¹⁵, K. Osterberg¹⁵, A. Ouraou⁴⁰, A. Oyanguren⁵⁰, M. Paganoni²⁹, S. Paiano⁵, P. Palacios²³, H. Palka¹⁸, D. Papadopoulou³², L. Pape⁸, C. Parkes²⁴, F. Parodi¹³, U. Parzefall⁸, A. Passeri³⁹, O. Passon⁵³, L. Peralta²², V. Perepelitsa⁵⁰, A. Perrotta⁵, A. Petrolini¹³, J. Piedra⁴¹, L. Pieri³⁹, F. Pierre⁴⁰, M. Pimenta²², E. Piotto⁸, T. Podobnik⁴³, V. Poireau⁸, E. Pol⁶, G. Polok¹⁸, P. Poropat⁴⁷, V. Pozdniakov¹⁶, N. Pukhaeva^{2,16}, A. Pullia²⁹, J. Rames¹², L. Ramler¹⁷, A. Read³³, P. Rebecchi⁸, J. Rehn¹⁷, D. Reid³¹, R. Reinhardt⁵³, P. Renton³⁵, F. Richard²⁰, J. Ridky¹², M. Rivero⁴¹, D. Rodriguez⁴¹, A. Romero⁴⁵, P. Ronchese³⁶, E. Rosenberg¹, P. Roudeau²⁰, T. Rovelli⁵, V. Ruhlmann-Kleider⁴⁰, D. Ryabtchikov⁴², A. Sadovsky¹⁶, L. Salmi¹⁵, J. Salt⁵⁰, A. Savoy-Navarro²⁵, U. Schwickerath⁸, A. Segar³⁵, R. Sekulin³⁷, M. Siebel⁵³, A. Sisakian¹⁶, G. Smadja²⁷, O. Smirnova²⁶, A. Sokolov⁴², A. Sopczak²¹, R. Sosnowski⁵², T. Spassov⁸, M. Stanitzki¹⁷, A. Stocchi²⁰, J. Strauss⁵¹, B. Stugu⁴, M. Szczekowski⁵², M. Szeptycka⁵², T. Szumlak¹⁹, T. Tabarelli²⁹, C. Taffard²³, F. Tegenfeldt⁴⁹, J. Timmermans³¹, L. Tkatchev¹⁶, M. Tobin²³, S. Todorovova¹², B. Tome²², A. Tonazzo²⁹, P. Tortosa⁵⁰, P. Travnicek¹², D. Treille⁸, G. Tristram⁷, M. Trochimczuk⁵², C. Troncon²⁸, M-L. Turluer⁴⁰, A. Tyapkin¹⁶, P. Tyapkin¹⁶, S. Tzamarias¹¹, V. Uvarov⁴², G. Valenti⁵, P. Van Dam³¹, J. Van Eldik⁸, A. Van Lysebetten², N. van Remortel², I. Van Vulpen⁸, G. Vegni²⁸, F. Veloso²², W. Venus³⁷, P. Verdier²⁷, V. Verzi³⁸, D. Vilanova⁴⁰, L. Vitale⁴⁷, V. Vrba¹², H. Wahlen⁵³, J. Washbrook²³, C. Weiser¹⁷, D. Wicke⁸, J. Wickens², G. Wilkinson³⁵, M. Winter⁹, M. Witek¹⁸, O. Yushchenko⁴², A. Zalewska¹⁸, P. Zalewski⁵², D. Zavrtanik⁴³, V. Zhuravlov¹⁶, I. Zimin¹⁶, A. Zintchenko¹⁶, M. Zupan¹¹

- ¹ Department of Physics and Astronomy, Iowa State University, Ames IA 50011-3160, USA
- ² Physics Department, Universiteit Antwerpen, Universiteitsplein 1, 2610 Antwerpen, Belgium and IIHE, ULB-VUB, Pleinlaan 2, 1050 Brussels, Belgium
and Faculté des Sciences, Univ. de l'Etat Mons, Av. Maistriau 19, 7000 Mons, Belgium
- ³ Physics Laboratory, University of Athens, Solonos Str. 104, 10680 Athens, Greece
- ⁴ Department of Physics, University of Bergen, Allégaten 55, 5007 Bergen, Norway
- ⁵ Dipartimento di Fisica, Università di Bologna and INFN, Via Irnerio 46, 40126 Bologna, Italy
- ⁶ Centro Brasileiro de Pesquisas Físicas, rua Xavier Sigaud 150, 22290 Rio de Janeiro, Brazil
and Depto. de Física, Pont. Univ. Católica, C.P. 38071, 22453 Rio de Janeiro, Brazil
and Inst. de Física, Univ. Estadual do Rio de Janeiro, rua São Francisco Xavier 524, Rio de Janeiro, Brazil
- ⁷ Collège de France, Lab. de Physique Corpusculaire, IN2P3-CNRS, 75231 Paris Cedex 05, France
- ⁸ CERN, 1211 Geneva 23, Switzerland
- ⁹ Institut de Recherches Subatomiques, IN2P3 - CNRS/ULP - BP20, 67037 Strasbourg Cedex, France
- ¹⁰ Now at DESY-Zeuthen, Platanenallee 6, 15735 Zeuthen, Germany
- ¹¹ Institute of Nuclear Physics, N.C.S.R. Demokritos, P.O. Box 60228, 15310 Athens, Greece
- ¹² FZU, Inst. of Phys. of the C.A.S. High Energy Physics Division, Na Slovance 2, 180 40, Praha 8, Czech Republic
- ¹³ Dipartimento di Fisica, Università di Genova and INFN, Via Dodecaneso 33, 16146 Genova, Italy
- ¹⁴ Institut des Sciences Nucléaires, IN2P3-CNRS, Université de Grenoble 1, 38026 Grenoble Cedex, France
- ¹⁵ Helsinki Institute of Physics, P.O. Box 64, 00014 University of Helsinki, Finland
- ¹⁶ Joint Institute for Nuclear Research, Dubna, Head Post Office, P.O. Box 79, 101 000 Moscow, Russian Federation
- ¹⁷ Institut für Experimentelle Kernphysik, Universität Karlsruhe, Postfach 6980, 76128 Karlsruhe, Germany
- ¹⁸ Institute of Nuclear Physics, Ul. Kawiory 26a, 30055 Krakow, Poland
- ¹⁹ Faculty of Physics and Nuclear Techniques, University of Mining and Metallurgy, 30055 Krakow, Poland
- ²⁰ Université de Paris-Sud, Lab. de l'Accélérateur Linéaire, IN2P3-CNRS, Bât. 200, 91405 Orsay Cedex, France
- ²¹ School of Physics and Chemistry, University of Lancaster, Lancaster LA1 4YB, UK
- ²² LIP, IST, FCUL - Av. Elias Garcia, 14-1^o, 1000 Lisboa Codex, Portugal
- ²³ Department of Physics, University of Liverpool, P.O. Box 147, Liverpool L69 3BX, UK
- ²⁴ Dept. of Physics and Astronomy, Kelvin Building, University of Glasgow, Glasgow G12 8QQ
- ²⁵ LPNHE, IN2P3-CNRS, Univ. Paris VI et VII, Tour 33 (RdC), 4 place Jussieu, 75252 Paris Cedex 05, France
- ²⁶ Department of Physics, University of Lund, Sölvegatan 14, 223 63 Lund, Sweden
- ²⁷ Université Claude Bernard de Lyon, IPNL, IN2P3-CNRS, 69622 Villeurbanne Cedex, France
- ²⁸ Dipartimento di Fisica, Università di Milano and INFN-MILANO, Via Celoria 16, 20133 Milan, Italy
- ²⁹ Dipartimento di Fisica, Univ. di Milano-Bicocca and INFN-MILANO, Piazza della Scienza 2, 20126 Milan, Italy
- ³⁰ IPNP of MFF, Charles Univ., Areal MFF, V Holesovickach 2, 180 00, Praha 8, Czech Republic
- ³¹ NIKHEF, Postbus 41882, 1009 DB Amsterdam, The Netherlands
- ³² National Technical University, Physics Department, Zografou Campus, 15773 Athens, Greece
- ³³ Physics Department, University of Oslo, Blindern, 0316 Oslo, Norway
- ³⁴ Dpto. Física, Univ. Oviedo, Avda. Calvo Sotelo s/n, 33007 Oviedo, Spain
- ³⁵ Department of Physics, University of Oxford, Keble Road, Oxford OX1 3RH, UK
- ³⁶ Dipartimento di Fisica, Università di Padova and INFN, Via Marzolo 8, 35131 Padua, Italy
- ³⁷ Rutherford Appleton Laboratory, Chilton, Didcot OX11 0QX, UK
- ³⁸ Dipartimento di Fisica, Università di Roma II and INFN, Tor Vergata, 00173 Rome, Italy
- ³⁹ Dipartimento di Fisica, Università di Roma III and INFN, Via della Vasca Navale 84, 00146 Rome, Italy
- ⁴⁰ DAPNIA/Service de Physique des Particules, CEA-Saclay, 91191 Gif-sur-Yvette Cedex, France
- ⁴¹ Instituto de Física de Cantabria (CSIC-UC), Avda. los Castros s/n, 39006 Santander, Spain
- ⁴² Inst. for High Energy Physics, Serpukov P.O. Box 35, Protvino, (Moscow Region), Russian Federation
- ⁴³ J. Stefan Institute, Jamova 39, 1000 Ljubljana, Slovenia and Laboratory for Astroparticle Physics, Nova Gorica Polytechnic, Kostanjeviska 16a, 5000 Nova Gorica, Slovenia,
and Department of Physics, University of Ljubljana, 1000 Ljubljana, Slovenia
- ⁴⁴ Fysikum, Stockholm University, Box 6730, 113 85 Stockholm, Sweden
- ⁴⁵ Dipartimento di Fisica Sperimentale, Università di Torino and INFN, Via P. Giuria 1, 10125 Turin, Italy
- ⁴⁶ INFN, Sezione di Torino, and Dipartimento di Fisica Teorica, Università di Torino, Via P. Giuria 1, 10125 Turin, Italy
- ⁴⁷ Dipartimento di Fisica, Università di Trieste and INFN, Via A. Valerio 2, 34127 Trieste, Italy
and Istituto di Fisica, Università di Udine, 33100 Udine, Italy
- ⁴⁸ Univ. Federal do Rio de Janeiro, C.P. 68528 Cidade Univ., Ilha do Fundão 21945-970 Rio de Janeiro, Brazil
- ⁴⁹ Department of Radiation Sciences, University of Uppsala, P.O. Box 535, 751 21 Uppsala, Sweden
- ⁵⁰ IFIC, Valencia-CSIC, and D.F.A.M.N., U. de Valencia, Avda. Dr. Moliner 50, 46100 Burjassot (Valencia), Spain
- ⁵¹ Institut für Hochenergiephysik, Österr. Akad. d. Wissensch., Nikolsdorfergasse 18, 1050 Vienna, Austria
- ⁵² Inst. Nuclear Studies and University of Warsaw, Ul. Hoza 69, 00681 Warsaw, Poland
- ⁵³ Fachbereich Physik, University of Wuppertal, Postfach 100 127, 42097 Wuppertal, Germany
- † deceased

Received: 18 September 2003 / Revised version: 26 March 2004 /

Published online: 11 May 2004 – © Springer-Verlag / Società Italiana di Fisica 2004

Abstract. Tau-pair production in the process $e^+e^- \rightarrow e^+e^-\tau^+\tau^-$ was studied using data collected by the DELPHI experiment at LEP2 during the years 1997 – 2000. The corresponding integrated luminosity is 650 pb^{-1} . The values of the cross-section obtained are found to be in agreement with QED predictions. Limits on the anomalous magnetic and electric dipole moments of the tau lepton are deduced.

1 Introduction

This paper presents a study of tau pair production in photon-photon collisions using the data collected by the DELPHI detector at LEP in the period from 1997 to 2000 (LEP2) at collision energy \sqrt{s} between 183 and 208 GeV. The total integrated luminosity used in the analysis is 650 pb^{-1} . At LEP this process was first observed by the OPAL collaboration [1] and subsequently studied by the L3 collaboration [2].

The final state $e^+e^-\tau^+\tau^-$ can be produced via a set of Feynman diagrams. In this paper we present the cross-section measurement for the contribution of the so-called multiperipheral graph (Fig. 1) which corresponds to collisions of two virtual photons. The same final states produced via other diagrams (less than 1% of the cross-section) are considered as a background.

The study of the reaction $e^+e^- \rightarrow e^+e^-\tau^+\tau^-$ explores two fundamental problems. First of all it provides a deep test of QED at the level of the fourth order in α . Furthermore, the $\gamma\tau\tau$ vertex is sensitive to the anomalous electromagnetic couplings of the tau lepton. Since the multiperipheral $e^+e^- \rightarrow e^+e^-\tau^+\tau^-$ process diagram contains two such vertices, the anomalous magnetic and electric dipole moments can be extracted by comparing the measured cross-section with QED expectations.

The rest of the paper gives a detailed description of the $e^+e^- \rightarrow e^+e^-\tau^+\tau^-$ cross-section measurement, including tau-pair selection, background estimation, selection and trigger efficiency calculation and systematic error estimation. In the last part of the paper the measured cross-sections are used to derive limits on the anomalous electromagnetic moments of the tau lepton.




Fig. 1. The dominant diagram for the reaction $e^+e^- \rightarrow e^+e^-\tau^+\tau^-$

2 Monte Carlo simulation

The signal process was simulated using the Berends, Daverveldt and Kleiss generator RADCOR (BDKRC) [3], which calculates the cross-section for the multiperipheral diagram with radiative corrections on the electron and positron lines. The following signal definition was used: the invariant mass of tau pairs had to be less than $40 \text{ GeV}/c^2$; both beam particles had to be scattered by less than 10 degrees; and at least one of them had to be scattered by less than 2 degrees. With these restrictions the accepted cross-section was $1.44 \pm 0.04\%$ lower than the total cross-section predicted by BDKRC (which is about 450 pb at LEP2 energies) for the unrestricted phase space. The τ decay was simulated by the TAUOLA package [4], which includes photon radiation from the decay products. The BDKRC generator was also used to estimate the background coming from the process $e^+e^- \rightarrow e^+e^-\mu^+\mu^-$.

To simulate the $e^+e^- \rightarrow e^+e^-e^+e^-$ background, the Berends, Daverveldt and Kleiss generator DIAG36 (BDK) [5] was used. Hadron production in two photon collisions was simulated by PYTHIA 6.1 [6]. Non-multiperipheral four-fermion processes (such as WW, ZZ, Zee and others) were simulated by WPHACT [7].

The generated events were passed through the full simulation program of the DELPHI detector and were reconstructed with the same program as for the real data [8].

3 Event selection

In most events produced by two-photon collisions both beam particles scatter at small angles and remain undetectable inside the beam pipe. Therefore only the decay products of the tau leptons can be seen in the detector. To suppress background, only one-prong decay channels with one tau decaying into an electron and the other into a non-electron (hadron or muon) were considered. The analysis was based entirely on the measured tracks of charged products of tau decays; the neutral particles from tau decays were ignored in this analysis.

To select runs with good performance of the sub-detectors [8–9], only runs with the Time Projection Chamber (TPC), the Forward Chambers (FCA, FCB) and one of the additional barrel tracking detectors (ID or VD) fully operational were retained. Table 1 presents the luminosities used in the analysis, luminosity-weighted centre-of-mass energies and energy ranges.

The event selection procedure was divided into two steps. The first step (preselection) selected a sample of two-photon events with two good tracks which were not

Table 1. The integrated luminosities, mean collision energies and collision energy ranges

	1997	1998	1999	2000
Luminosity, pb ⁻¹	52.3	152.6	224.2	217.5
$\langle E_{cm} \rangle$, GeV	182.7	188.7	197.6	206.3
Energy range, GeV	182.7	188.7	195.5–201.5	204.5–208.0

back-to-back in the plane perpendicular to the beam axis. A track was considered as good if the momentum derived from its curvature was greater than 100 MeV/c, momentum error better than 100%, polar angle θ between 20° and 160°, and impact parameter with respect to the interaction point below 10 cm along the z -axis¹ and 5 cm in the $r - \phi$ plane.

The following cuts were applied in this first selection step:

- There had to be exactly two good tracks from particles with opposite charges, at least one of them having momentum greater than 300 MeV/c.
- To suppress background from fermion pair production, the total energy of two charged particles had to be less than 30 GeV.

- To enrich the sample with $e^+e^- \rightarrow e^+e^-\tau^+\tau^-$ events, the acoplanarity of two tracks² had to be greater than 0.5° and their resultant transverse momentum greater than 500 MeV/c.
- To select events with a high trigger efficiency, the transverse energy, defined by

$$E_t = E_1 \sin \theta_1 + E_2 \sin \theta_2,$$

where E_1 and E_2 are the energies of the two charged particles and θ_1 and θ_2 are their polar angles, had to be greater than 2 GeV.

- In the year 2000, the operation of one of the twelve TPC sectors was unstable and the dE/dx measurement vital for this analysis was poor, so events with at least one track in or near (closer than 10° in ϕ) to this TPC sector in 2000 were rejected.
- Finally, to ensure the transverse momentum balance of $\gamma\gamma$ system, single and double tagged events were rejected by requiring that no energy deposition in the forward electromagnetic calorimeters (STIC or FEMC) exceeded 60% of the beam energy.




Fig. 2. The distributions of invariant mass, total energy, transverse energy and transverse momentum of the pair of charged particles. Preselection cuts are applied. The points are 1999 data, the open histogram is the simulation of background processes and the shaded histogram is the simulation of $e^+e^- \rightarrow e^+e^-\tau^+\tau^-$ events. The simulation is not corrected for the trigger efficiency

¹ The DELPHI coordinate system has the z -axis aligned along the electron beam direction, the x -axis pointing toward the centre of LEP and the y -axis vertical. r is the radius in

the (x, y) plane. The polar angle θ is measured with respect to the z -axis and the azimuthal angle ϕ is about z .

² Acoplanarity is defined as $180^\circ - |\phi_2 - \phi_1|$.

The last cut suppressed the events with highly virtual photons. About 90% of events passing this cut had the momentum transfer $-q^2$ less than $1 \text{ GeV}^2/c^2$. After applying the cuts described above, the predicted event composition in the preselected sample was as follows (1999 data):

$e^+e^- \rightarrow e^+e^-e^+e^-$	41%
$e^+e^- \rightarrow e^+e^-\mu^+\mu^-$	47%
$e^+e^- \rightarrow e^+e^-\tau^+\tau^-$	8%
$e^+e^- \rightarrow e^+e^-q\bar{q}$	3%
$e^+e^- \rightarrow \tau^+\tau^-$	1%

The fraction of other events was less than 1%. The efficiency of the preselection for $e^+e^- \rightarrow e^+e^-\tau^+\tau^-$ events was of the order of 5%, the largest suppression of the signal coming from the requirement of exactly two good tracks seen in the detector (about a factor of 4) and from the cut on E_t (about factor of 2). Figure 2 shows the comparison between data and simulation of the distributions of invariant mass, total energy, total transverse energy and total transverse momentum of the pair of charged particles. The $e^+e^- \rightarrow e^+e^-\tau^+\tau^-$ events are shown by the shaded histogram. The Monte Carlo is normalised to the luminosity of the real data. The data deficit is mainly due to the trigger inefficiency which is corrected at the later stages of analysis.

In the final step of the selection, the event was retained if one of the charged particles was identified as an electron and the other as a non-electron. This step was based on the TPC measurement of the dE/dx pulls for the muon, electron, kaon and proton hypotheses. The dE/dx pull for a specific particle hypothesis is defined as the ratio

$$\Pi_X = \frac{(dE/dx)_{\text{meas}} - (dE/dx)_{\text{exp}}}{\sigma_{dE/dx}}, \quad (1)$$

where $(dE/dx)_{\text{exp}}$ is the value expected for the particle X with given momentum and $\sigma_{dE/dx}$ is the error of the measured energy loss $(dE/dx)_{\text{meas}}$. To check the dE/dx calibration, test samples of $e^+e^- \rightarrow e^+e^-\mu^+\mu^-$ and $e^+e^- \rightarrow e^+e^-e^+e^-$ events were picked out from the preselected sample. A small angular dependence of the dE/dx measurements was found as well as some disagreement between data and simulation. Corrections which were functions of azimuthal and polar angle were applied to the measured dE/dx values. Residual disagreement was removed by scaling and smearing the specific energy loss measurement in the simulated events. Independent calibrations of real and simulated data were performed for each year of data taking analysed. The efficiency to measure dE/dx is discussed later in the paper.

With corrected dE/dx information, a track was identified as an electron if $\Pi_\mu > 3$ and as a non-electron if $\Pi_e < -3$. Figure 3 illustrates the particle identification cuts. The distributions of the pulls for the electron and muon hypotheses are shown for the 1999 real data and simulation. Each distribution is shown after applying all selection cuts except the cut on the variable shown. The shaded histograms show the signal.

Table 2. Efficiencies (%): preselection, final step of selection and overall efficiency

	1997	1998	1999	2000
preselection	5.39	5.37	5.38	3.85
selection	17.3	16.4	16.4	16.1
overall	0.93	0.88	0.88	0.62

A considerable amount of kaon and proton background from $e^+e^- \rightarrow e^+e^-q\bar{q}$ events remained after the cuts on the pulls for the muon and electron hypotheses. Figure 4 (left) shows the specific energy loss for electron candidates plotted versus the momentum of the particle. Proton and kaon bands are clearly visible. To remove the kaon and proton background, the electron selection was tightened. The dE/dx for the electron candidate had to not exceed 1.9 times the minimum ionisation, and the pulls for the proton and kaon hypotheses for the electron candidate both had to be outside the $\pm 1.5\sigma$ interval: $|\Pi_K| > 1.5$ and $|\Pi_p| > 1.5$. Figure 4 (right) shows the distribution of the pull for the proton hypothesis with all selection cuts applied except the cut on the variable shown. The hatched histogram shows the background from $e^+e^- \rightarrow e^+e^-q\bar{q}$ events, the shaded histogram shows the rest of the background. The cuts on this variable are indicated by arrows.

Table 2 summarises the efficiency of the preselection, final step of selection and overall selection efficiency. The drop in the preselection in 2000 is caused by the removal of events in or near the unstable TPC sector. The uncertainties of the selection efficiency determination are discussed later in the paper.

In total 2390 candidate events were selected. Figure 5 compares the distributions of electron and non-electron candidate momenta for selected events to the simulation prediction for the combined 1997–2000 data. Figure 6 shows the visible invariant mass distribution for selected events for the same data sample. Trigger efficiency is taken into account in these distributions (see below).

3.1 Trigger efficiency

The low momenta of the τ decay products in the process $e^+e^- \rightarrow e^+e^-\tau^+\tau^-$ and the presence of only two tracks in the event could make the probability of triggering on such an event considerably below 100%. The determination of the trigger efficiency is therefore important in this analysis.

The trigger efficiency was estimated from the subsamples of selected events using the fact that an event can be detected by different components of the DELPHI trigger system [10]. Trigger subcomponents of the tracking system were combined into barrel and end-cap triggers. For events with one track in the barrel and the other in the end-cap, the number of events detected by the barrel trigger (N_B), by the end-cap trigger (N_E), and by both triggers (N_{BE}) were counted using the decision functions of the trigger. The barrel and end-cap single track trigger efficiencies were calculated, for electrons and non-electrons separately, by




Fig. 3. Distribution of the dE/dx pull for the electron hypothesis (left) and for the muon hypothesis (right) with all selection cuts applied except the cut on the variable shown. Points are 1999 data, the open histogram is the background, and the shaded histogram shows the $e^+e^- \rightarrow e^+e^-\tau^+\tau^-$ signal events. The simulation is not corrected for the trigger efficiency




Fig. 4. Left: specific energy loss as a function of particle momentum for electron candidates after preselection cuts and the cuts on the electron and muon pulls. The horizontal line shows the first cut against kaons and protons. The points are 1999 data. Right: distribution of dE/dx pull for proton hypothesis for electron candidates. The hatched histogram is the background from $e^+e^- \rightarrow e^+e^-q\bar{q}$ events and the shaded histogram is the rest of the background. The cuts against protons are indicated by arrows. All other selection cuts are applied. The points are 1999 data




Fig. 5. Momentum distribution for electron candidates (left) and non-electron candidates (right) for selected events from 1997–2000 data. The distributions of simulated events are corrected for trigger efficiency. The dip in the electron momentum distribution near 1 GeV/c is caused by the cut against protons: the electron and proton dE/dx are equal in this region of momentum

the formulae:

$$\varepsilon_{\text{Barrel}} = \frac{N_{\text{BE}}}{N_{\text{E}}}; \quad \varepsilon_{\text{end-cap}} = \frac{N_{\text{BE}}}{N_{\text{B}}}. \quad (2)$$

Finally, the efficiency of the DELPHI calorimetric trigger to the whole event was estimated from the events triggered by any of the tracking detectors using a similar technique. The results of the trigger efficiency calculation are summarised in Table 3. The track pair trigger efficiency was calculated from the “OR” of the single track efficiencies using the ratio

of the barrel and forward tracks predicted by simulation. The tau pair trigger efficiency was calculated as “OR” of the tracking and calorimetric triggers.

3.2 Efficiency of the dE/dx measurement

Both tracks of the selected event had to have specific energy loss measurements. An imperfect detector simulation can lead to a discrepancy in the dE/dx measurement effi-

Table 3. Summary of the trigger efficiency measurements (%)

	1997	1998	1999	2000
Barrel track				
electron	71.4 ± 17.1	94.4 ± 5.4	84.6 ± 7.1	92.3 ± 7.4
non-electron	$100^{+0}_{-17.9}$	85.2 ± 6.8	85.0 ± 8.0	78.6 ± 11.0
End-cap track				
electron	26.3 ± 10.1	36.5 ± 6.1	21.5 ± 4.6	22.0 ± 6.9
non-electron	31.3 ± 11.6	30.4 ± 6.1	25.9 ± 4.8	23.1 ± 5.8
Track pair	$94.5^{+5.5}_{-7.1}$	95.5 ± 2.7	93.3 ± 3.5	93.5 ± 4.1
Calorimetry	6.7 ± 1.9	8.6 ± 1.2	7.1 ± 0.9	7.7 ± 1.1
Tau pair	$94.9^{+5.1}_{-6.6}$	95.9 ± 2.5	93.8 ± 3.3	94.0 ± 3.8




Fig. 6. Visible invariant mass distribution for selected events for combined 1997–2000 data. The distribution of simulated events is corrected for trigger efficiency. The mass was calculated using all detected charged particles and photons. The simulation was corrected for the trigger efficiency

ciency for good tracks in real and simulated events. To take into account this possible disagreement, the efficiency for a good track to have a dE/dx measurement (to be a “good TPC track”) was calculated for $e^+e^- \rightarrow e^+e^-e^+e^-$ and $e^+e^- \rightarrow e^+e^-\mu^+\mu^-$ samples extracted from preselected events (efficiencies to measure dE/dx of pions and muons were assumed to be equal). Muon events were selected by requiring at least one track to be identified by the muon chambers and electron events were selected using information from the DELPHI RICH detectors. For muon and electron events the efficiency to be a “good TPC track” was determined from the ratio

$$\varepsilon_{dE/dx}^2 = \frac{N_{dE/dx}}{N_{tot}} \quad (3)$$

Table 4. Summary of “good TPC track” efficiency estimations. These efficiencies are already included in the total efficiency in Table 2

	1997	1998	1999	2000
Efficiency in data, %	82.9 ± 1.0	82.6 ± 0.6	82.4 ± 0.5	83.5 ± 0.6
Efficiency in MC, %	82.3 ± 0.4	82.5 ± 0.2	82.3 ± 0.1	84.6 ± 0.2

where $N_{dE/dx}$ is the number of events with both tracks having a dE/dx measurement and N_{tot} was the total number of selected events in the given sample. The tau-pair efficiency was estimated as the product of the single track efficiencies for muon and electron. The tau-pair efficiencies derived from $e^+e^- \rightarrow e^+e^-\mu^+\mu^-$ and $e^+e^- \rightarrow e^+e^-e^+e^-$ events for data and Monte Carlo are presented in Table 4 and were used for selection efficiency correction and for systematic error estimation. The selection efficiency was multiplied by the factor $\frac{\varepsilon_{dE/dx}(\text{data})}{\varepsilon_{dE/dx}(\text{MC})}$ and half of the correction was included into the systematic error together with the uncertainties from the test sample statistics.

3.3 Residual background

Several sources of background for $e^+e^- \rightarrow e^+e^-\tau^+\tau^-$ events have been considered:

- The background from $e^+e^- \rightarrow e^+e^-q\bar{q}$, mainly from protons and kaons selected due to the tails of the dE/dx pulls for the proton and kaon hypotheses;
- The background from $e^+e^- \rightarrow e^+e^-e^+e^-$ and $e^+e^- \rightarrow e^+e^-\mu^+\mu^-$ events due to the tails of the distributions of the dE/dx pulls for the electron and muon hypotheses;
- Background due to other four-fermion processes: non-multiperipheral diagrams (including $e^+e^-\tau^+\tau^-$ final states) and multiperipheral process $e^+e^- \rightarrow e^+e^-\tau^+\tau^-$ which does not satisfy signal definition;
- The process $e^+e^- \rightarrow \tau^+\tau^-$ (background from other fermion pair production processes was found to be negligible).

The background fractions for the main background sources and their uncertainties are summarised in Table 5. The contribution from other background sources was negligible. The theoretical precision of $e^+e^- \rightarrow e^+e^-q\bar{q}$ generation by PYTHIA is not well known. Therefore it was

Table 5. Summary of background fractions. The numbers are the expected fractions (%) of the specified backgrounds in the selected sample. Errors are statistical errors of the simulated samples and theoretical uncertainties of the Monte Carlo generators added in quadrature

Channel	1997	1998	1999	2000
$e^+e^- \rightarrow e^+e^-q\bar{q}$	4.3 ± 1.5	3.2 ± 0.7	3.3 ± 0.8	3.2 ± 0.8
$e^+e^- \rightarrow e^+e^-e^+e^-$	2.7 ± 0.2	3.4 ± 0.1	4.0 ± 0.1	2.4 ± 0.1
$e^+e^- \rightarrow e^+e^-\mu\mu$	2.9 ± 0.1	5.0 ± 0.1	2.4 ± 0.1	3.8 ± 0.1
other 4-fermion	1.5 ± 0.3	1.5 ± 0.3	1.2 ± 0.2	1.2 ± 0.2
$e^+e^- \rightarrow \tau\tau$	0.69 ± 0.01	0.55 ± 0.01	0.47 ± 0.01	0.40 ± 0.01
Total	12.1 ± 1.5	13.6 ± 0.8	11.4 ± 0.8	11.0 ± 0.8

estimated from the real data by inverting the dE/dx cut on the electron candidate: $dE/dx > 1.9$ M.I.P. instead of $dE/dx < 1.9$ M.I.P. After comparing these test samples enriched with $e^+e^- \rightarrow e^+e^-q\bar{q}$ events with the simulation, an error of 20% was ascribed to the $e^+e^- \rightarrow e^+e^-q\bar{q}$ event generator.

4 Systematic error estimation

The following sources of systematic error on the cross-section measurement were considered: uncertainties of selection and trigger efficiencies and uncertainty of background level. Track selection, event selection and the statistical error of the simulated samples were taken into account in the calculation of the uncertainty in the selection efficiency.

The systematic error arising from track selection was estimated in the following way. Each cut of the track selection was varied typically by the size of the resolution of the corresponding variable from its nominal value in both directions. The corresponding change of the cross-section Δ was compared to the value of the expected statistical fluctuation σ due to the non-identical event sample. If the value Δ was less than σ , no systematic error was ascribed to the corresponding cut; in the opposite case the value of $\sqrt{\Delta^2 - \sigma^2}$ was included in the systematic error. The systematic error arising from varying the event selection cuts was estimated in a similar way.

To calculate the systematic error due to the angular corrections applied to the dE/dx measurements, the dE/dx correction functions were varied by the uncertainty of each of their parameters and the analysis chain was repeated. The variation of the measured cross-section was added to the systematic error. The systematic errors corresponding to scaling and smearing the pulls were calculated similarly.

The systematic errors associated with track selection cuts, event selection cuts and dE/dx corrections are summarised in Table 6. The numbers are given for 1999 data (for other years uncertainties of most of the sources scale approximately as inverse square root of the statistics). Additional contributions to the selection efficiency uncertainty also presented in Table 6 come from the statistical error of the Monte Carlo sample and the selection efficiency correction described in Sect. 3.2.

Table 6. Systematic errors for 1999 data coming from track selection, event selection, dE/dx corrections, simulated samples statistics and “good TPC track” correction

syst. error source	value, %
track selection cuts	
R_{imp}	0.7
Z_{imp}	1.1
$\delta p/p$	0.7
event selection cuts	
Π_e	0.3
Π_μ	0.3
acoplanarity	0.6
dE/dx corrections	
$\Pi_e \theta$	1.0
$\Pi_e \phi$	0.9
$\Pi_\mu \theta$	1.0
$\Pi_\mu \phi$	1.0
scaling	0.7
smearing	0.6
MC statistics	0.8
“Good TPC track”	
correction	0.6
Total	3.0

The largest contribution to the systematic error is given by the uncertainty of the trigger efficiency determination, dominated by the statistics of the real data events, see Sect. 3.1 and Table 3. An additional contribution arises because the trigger efficiency for background events, assumed to be equal to that of the signal, may be different. A conservative estimate of this uncertainty was obtained by changing the trigger efficiency for background upwards to 100% and downwards by the same amount.

The systematic error due to residual background includes the simulated sample statistical uncertainty and the theoretical uncertainty of the Monte Carlo generators, mainly for the $e^+e^- \rightarrow e^+e^-q\bar{q}$ process, see Table 5.

The sources of selection efficiency uncertainty are described in detail in Table 6.

The sources of systematic uncertainty are summarised in Table 7. Total systematic errors calculated as the sum in

Table 7. Relative systematic errors on cross-section (in %)

	1997	1998	1999	2000
Trigger eff.	7.0	2.7	3.6	4.5
Selection eff.	5.1	3.2	3.0	3.0
Background	1.7	0.9	0.9	0.9
Luminosity	0.6	0.6	0.6	0.6
Total	8.9	4.3	4.7	5.4

quadrature of all described components are also presented in Table 7. The following uncertainties were assumed to be fully correlated between different years: generator theoretical error; trigger efficiency for background; and uncertainties estimated from variation of track and event selection cuts. Systematic errors from other sources were treated as uncorrelated.

5 Results of the cross-section measurement

The cross-sections were computed using the formula

$$\sigma = \frac{N_{\text{obs}} - N_{\text{bg}}}{\varepsilon_{\text{sel}}\varepsilon_{\text{trig}}\mathcal{L}} \quad (4)$$

where N_{obs} is the number of observed events, N_{bg} is the expected number of background events in the assumption that background events have the same trigger efficiency as signal events, ε_{sel} is the selection efficiency, $\varepsilon_{\text{trig}}$ is the trigger efficiency and \mathcal{L} is the integrated luminosity.

The numbers of observed and expected events, the measured cross-sections and the cross-sections from the BDKRC Monte Carlo simulation together with their ratios are presented in Table 8. The predicted number of events was calculated from the signal and background simulation, taking into account trigger efficiency and corrections to the dE/dx efficiency. Agreement was found between the measurements and the Standard Model (SM) predictions calculated by BDKRC. The ratio of observed and predicted cross-sections was averaged over all LEP2 data, taking into account correlations of systematic errors. The result was found to be 0.96 ± 0.04 . The average LEP2 cross-section is 429 ± 17 pb corresponding to the luminosity-weighted mean centre-of-mass energy of 197.1 GeV. The cross-section predicted at this energy by BDKRC is 447.7 ± 0.3 pb.

Table 8. The numbers of observed and expected events, measured cross-sections, QED predictions and their ratios. The first error on the measured cross-sections is statistical, the second is systematic

Year	Observed	Expected	σ_{meas} , pb	σ_{MC} , pb	$\sigma_{\text{meas}}/\sigma_{\text{MC}}$
1997	211	224 ± 18	$401 \pm 32 \pm 36$	428.2 ± 0.5	0.94 ± 0.11
1998	629	652 ± 24	$419 \pm 19 \pm 18$	436.7 ± 0.5	0.96 ± 0.06
1999	909	937 ± 39	$436 \pm 16 \pm 21$	448.5 ± 0.5	0.97 ± 0.06
2000	641	665 ± 32	$443 \pm 20 \pm 24$	459.4 ± 0.5	0.97 ± 0.07

6 Determination of anomalous magnetic and electric dipole moments

In the Standard Model, leptons are considered as point-like objects. Therefore the observation of a deviation of the magnetic or electric dipole moments of the leptons from their SM values would open a window onto the physics beyond the SM. The anomalous magnetic moments of the electron [11] and muon [12] are known with high precision, but the short life-time of the tau-lepton does not allow measurement of its anomalous moments with similar precision by a spin precession method.

The generalised form of the $\tau\tau\gamma$ vertex can be parametrised as follows:

$$\begin{aligned} & -i\bar{e}\bar{u}(p') \\ & \times \left\{ F_1(q^2)\gamma^\mu + iF_2(q^2)\sigma^{\mu\nu}\frac{q_\nu}{2m_\tau} + F_3(q^2)\gamma^5\sigma^{\mu\nu}\frac{q_\nu}{2m_\tau} \right\} \\ & \times u(p)\epsilon_\mu(q) \end{aligned} \quad (5)$$

where $\epsilon_\mu(q)$ is the polarization vector of the photon with momentum q . The form factor F_1 describes the distribution of electric charge and $e_\tau = eF_1(0)$, while F_2 and F_3 are form factors related to the anomalous magnetic moment a_τ and electric dipole moment d_τ :

$$a_\tau \equiv \frac{g_\tau - 2}{2} = F_2(0) \quad (6)$$

and

$$F_3(0) = -\frac{2m_\tau d_\tau}{e_\tau} \quad (7)$$

In the SM at tree level, $a_\tau = 0$ and $d_\tau = 0$. Accounting for loop diagrams gives a non-zero value to $a_\tau = 11773(3) \cdot 10^{-7}$ [13], while a non-zero value of d_τ is forbidden by both T invariance and P invariance.

The values of a_τ and d_τ have been measured by several groups. The L3 and OPAL collaborations [14, 15] studied radiative $Z \rightarrow \tau\tau\gamma$ events and set the following 95 % CL limits on the values of the anomalous magnetic and electric dipole moments:

$$\begin{aligned} & -0.052 < a_\tau < 0.058 \quad \text{and} \quad |d_\tau| < 3.1 \quad (10^{-16} e \cdot \text{cm}) \quad (\text{L3}), \\ & -0.068 < a_\tau < 0.065 \quad \text{and} \quad |d_\tau| < 3.7 \quad (10^{-16} e \cdot \text{cm}) \\ & \hspace{15em} (\text{OPAL}). \end{aligned}$$




Fig. 7. Total cross-section change as a function of anomalous magnetic moment and as a function of electric dipole moment

The best limit so far on d_τ was obtained by BELLE [16]:

$$\begin{aligned} -0.22 < \Re e(d_\tau) < 0.45 & \quad (10^{-16} e \cdot \text{cm}), \\ -0.25 < \Im m(d_\tau) < 0.08 & \quad (10^{-16} e \cdot \text{cm}). \end{aligned}$$

Other limits on a_τ and d_τ can be found in [17].

6.1 Limits from this analysis

Here we present the study of the anomalous magnetic and electric dipole moments of the tau lepton based on the analysis of the $e^+e^- \rightarrow e^+e^-\tau^+\tau^-$ cross-section. The study of anomalous couplings of tau leptons to photons at LEP in this channel was proposed in [18].

To model the contribution of non-SM anomalous magnetic and dipole moments we use the calculation by Cornet and Illana [19]. The calculation is based on computation of the matrix element of the process $\gamma\gamma \rightarrow \tau^+\tau^-$ in leading order of QED and its translation to the cross-section of the $e^+e^- \rightarrow e^+e^-\tau^+\tau^-$ process using the Equivalent Photon Approximation (EPA) [20]. The EPA parameter $(-q^2)_{\text{max}}$ (the upper limit of the integration over 4-momenta of the emitted photon) was chosen such that the total cross-section predicted by EPA (with SM values of anomalous electromagnetic moments) agreed with BDKRC calculation. According to the calculations [19] each of the anomalous terms of (5) would mainly modify the rate of tau pair production in the barrel region of the detector where the experimental selection has largest efficiency. This leads to a larger selection efficiency for the anomalous term contribution, improving in principle the limits obtained on anomalous moments. However in this paper we conservatively assumed that the standard and anomalous contributions have the same selection efficiency.

Figure 7 shows how the total cross-section changes as a function of the anomalous magnetic moment and as a function of the electric dipole moment. The three lines on each plot represent the calculation with $\sqrt{s} = 182.7, 195.5$ and 205.0 GeV. Increasing the collision energy slowly increases both non-SM contributions. However, increasing the magnitude of the anomalous magnetic moment can

either increase or decrease the cross-section while increasing that of the electric dipole moment tends only to increase the cross-section.

To compare the experimentally measured values of the cross-sections to the non-SM calculation, they were first converted from accepted to the total cross-sections, taking into account the 1.44% difference due to the signal definition (see Sect. 2). The validity of applying SM conversion factors is supported by the fact that the measured cross-sections are in good agreement with the SM prediction, which guarantees the smallness of the non-SM contribution, and by the fact that the correction itself is small.

Fits to the cross-sections measured in 1997, 1998, 1999 and 2000 were performed taking a_τ and d_τ as parameters. When fitting for a_τ , the value of d_τ was set to its SM value and *vice versa*. The errors on the cross section measurements were taken as the statistical and systematic errors added in quadrature.

To quote the obtained limits we used the following convention:

$$\int_{-\infty}^L \exp(-\chi^2/2) da_\tau = \int_R^{\infty} \exp(-\chi^2/2) da_\tau = \frac{1 - CL}{2} \quad (8)$$

where CL is the desired confidence level and L and R are lower and upper limits. A similar definition was used for d_τ . We quote central values μ and errors σ for moments according to

$$\sigma = \frac{R - L}{2}, \quad \mu = \frac{R + L}{2}. \quad (9)$$

where R and L are calculated with 68.3% confidence level.

Figure 8 shows the χ^2 as a function of the anomalous magnetic moment and as a function of the electric dipole moment. The results of the fit are:

$$\begin{aligned} -0.052 < a_\tau < 0.013, & \quad 95\% \text{ CL}, \\ |d_\tau| < 3.7 \cdot 10^{-16} e \cdot \text{cm}, & \quad 95\% \text{ CL}. \end{aligned}$$

The limit on a_τ improves the current PDG limit [21] based on the L3 result [14].

Figure 9 shows the the measured cross-section, average LEP2 cross-section and SM expectation as a function of \sqrt{s} .




Fig. 8. χ^2 as a function of anomalous magnetic moment and as a function of electric dipole moment




Fig. 9. Measured cross-section (circles), average LEP2 cross-section (square) and SM expectation as a function of \sqrt{s} . The two bands show the cross-section variation allowed due to anomalous magnetic and electric dipole moments within 95% limits from this analysis

Two bands superimposed on the plot represent the allowed region for the cross-section variation due to anomalous magnetic and electric dipole moments. The results expressed in the form of central value and error are the following:

$$a_\tau = -0.018 \pm 0.017,$$

$$d_\tau = (0.0 \pm 2.0) \cdot 10^{-16} e \cdot \text{cm}.$$

7 Conclusion

We have studied the reaction $e^+e^- \rightarrow e^+e^-\tau^+\tau^-$ with the data collected with the DELPHI detector during LEP2 operation in the years 1997–2000. The average LEP2 cross-section was found to be 429 ± 17 pb compared to 447.7 pb expected from the Standard model. The measured/predicted ratio 0.96 ± 0.04 agrees with the QED prediction at the level of one standard deviation. The measured cross-sections were used to extract limits on the anomalous magnetic

and electric dipole moments of the tau lepton. The 95% CL limits obtained are

$$-0.052 < a_\tau < 0.013,$$

$$|d_\tau| < 3.7 \cdot 10^{-16} e \cdot \text{cm}.$$

Acknowledgements. We are greatly indebted to our technical collaborators, to the members of the CERN-SL Division for the excellent performance of the LEP collider, and to the funding agencies for their support in building and operating the DELPHI detector. We acknowledge in particular the support of Austrian Federal Ministry of Education, Science and Culture, GZ 616.364/2-III/2a/98, FNRS-FWO, Flanders Institute to encourage scientific and technological research in the industry (IWT), Federal Office for Scientific, Technical and Cultural affairs (OSTC), Belgium, FINEP, CNPq, CAPES, FUJB and FAPERJ, Brazil, Czech Ministry of Industry and Trade, GA CR 202/99/1362, Commission of the European Communities (DG XII), Direction des Sciences de la Matière, CEA, France, Bundesministerium für Bildung, Wissenschaft, Forschung und Technologie, Germany, General Secretariat for Research and Technology, Greece, National Science Foundation (NSF) and Foundation for Research on Matter (FOM), The Netherlands, Norwegian Research Council, State Committee for Scientific Research, Poland, SPUB-M/CERN/PO3/DZ296/2000, SPUB-M/CERN/PO3/DZ297/2000 and 2P03B 104 19 and 2P03B 69 23(2002–2004) JNICT–Junta Nacional de Investigação Científica e Tecnológica, Portugal, Vedecka grantova agentura MS SR, Slovakia, Nr. 95/5195/134, Ministry of Science and Technology of the Republic of Slovenia, CICYT, Spain, AEN99-0950 and AEN99-0761, The Swedish Natural Science Research Council, Particle Physics and Astronomy Research Council, UK, Department of Energy, USA, DE-FG02-01ER41155, EEC RTN contract HPRN-CT-00292-2002. We thank F.Cornet for providing the calculation of the cross-section of anomalous tau-pair production in two-photon collisions.

References

1. OPAL Collaboration, R. Akers et al., *Z. Phys. C* **60**, 593 (1993)
2. L3 Collaboration, M. Acciarri et al., *Phys. Lett. B* **407**, 341 (1997)

3. F.A. Berends, P.H. Daverveldt, R. Kleiss, *Comput. Phys. Commun.* **40**, 271 (1986)
4. S. Jadach, J. Kühn, Z. Was, *Comput. Phys. Commun.* **64**, 275 (1991)
5. F.A. Berends, P.H. Daverveldt, R. Kleiss, *Comput. Phys. Commun.* **40**, 285 (1986)
6. T. Sjöstrand et al., *Comput. Phys. Commun.* **135**, 228 (2001)
7. E. Accomando, A. Ballestrero, *Comput. Phys. Commun.* **99**, 270 (1997); E. Accomando, A. Ballestrero, E. Maina, *Comput. Phys. Commun.* **150**, 166 (2003)
8. DELPHI Collaboration, P. Abreu et al., *Nucl. Instr. Methods A* **378**, 57 (1996)
9. DELPHI Collaboration, P. Aarnio et al., *Nucl. Instr. Methods A* **303**, 233 (1991);
10. A. Augustinus et al., The DELPHI Trigger System at LEP2 Energies, CERN-EP/2002-086, accepted by *Nucl. Instr. Methods A*
11. P.J. Mohr, B.N. Taylor, *J. Phys. Chem. Ref. Data* **28**, 1713 (1999); P.J. Mohr, B.N. Taylor, *Rev. Mod. Phys.* **72**, 351 (2000)
12. Muon $g-2$ Collaboration, G.W. Bennett et al., *Phys. Rev. Lett.* **89**, 101804 (2002)
13. M.A. Samuel, G. Li, R. Mendel, *Phys. Rev. Lett.* **67**, 668 (1991); erratum *Phys. Rev. Lett.* **69**, 995 (1992)
14. L3 Collaboration, M. Acciarri et al., *Phys. Lett. B* **434**, 169 (1998)
15. OPAL Collaboration, K. Ackerstaff et al., *Phys. Lett. B* **431**, 188 (1998)
16. BELLE Collaboration, K. Inami et al., *Phys. Lett. B* **551**, 16 (2003)
17. G.A. Gonzalez-Sprinberg, A. Santamaria, J. Vidal, *Nucl. Phys. B* **582**, 3 (2000); R. Escribano, E. Massó, *Phys. Lett. B* **395**, 369 (1997); D.J. Silverman, G.L. Shaw, *Phys. Rev. D* **27**, 1196 (1983); F. del Aguila, M. Sher, *Phys. Lett. B* **252**, 116 (1990); ARGUS Collaboration, H. Albrecht et al., *Phys. Lett. B* **485**, 37 (2000)
18. F. Cornet, J. Illana, *Phys. Rev. D* **53**, 1181 (1996)
19. F. Cornet, private communication; J. Illana, Estudio de las propiedades electromagnéticas del boson W y del lepton tau en procesos de dos fotones, doctoral thesis (in Spanish), University of Granada, (1995)
20. V.M. Budnev et al., *Phys. Rep.* **15**, 181 (1975)
21. K. Hagiwara et al., *Phys. Rev. D* **66**, 010001 (2002)