


# P-odd and spin effects in HIC(@NICA)

NICA Roundtable V, JINR,  
Dubna, August 28 2010

---

Oleg Teryaev  
JINR



# Outline

---

NICA speciality – chemical potential+vorticity?

- Chiral Vortaic Effect – Calculations?! – discrete curl in transport model (E. Bratkovskaya et al), HF – D. Voskresensky et al
- CVE & neutron asymmetries @ NICA
- Bilinear current correlators in medium and dilepton angular distributions
- Polarization Core-corona model – relation to hyperon polarization in PP
- Other signals for (local) C/P violations – decays forbidden in vacuum

# Anomaly in medium – new external lines in VVA graph


- Gauge field  $\rightarrow$  velocity


- CME  $\rightarrow$  CVE

- Kharzeev,  
Zhitnitsky (07) –  
EM current

- Straightforward  
generalization:  
any (e.g. baryonic)

current – neutron asymmetries@NICA -  
Rogachevsky, Sorin, OT - Arxive 1006.1331 (hep-ph)





# Baryon charge with neutrons – (Generalized) Chiral Vortaic Effect

---


- Coupling:  $e_j A_\alpha J^\alpha \Rightarrow \mu_j V_\alpha J^\alpha$

- Current:  $J_e^\gamma = \frac{N_c}{4\pi^2 N_f} \varepsilon^{\gamma\beta\alpha\rho} \partial_\alpha V_\rho \partial_\beta (\theta \sum_j e_j \mu_j)$

- - Uniform chemical potentials:  $J_i^\nu = \frac{\sum_j g_{i(j)} \mu_j}{\sum_j e_j \mu_j} J_e^\nu$

- - Rapidly (and similarly) changing chemical potentials:


$$J_i^0 = \frac{|\vec{\nabla} \sum_j g_{i(j)} \mu_j|}{|\vec{\nabla} \sum_j e_j \mu_j|} J_e^0$$



# Comparing CME and CVE

---

- Orbital Angular Momentum and magnetic moment are proportional – Larmor theorem
- Ideal fluid - Circulation conservation
- Same scale effect as that of magnetic field ( $eH/(\mu \text{ curl } v) \sim 1$ )
- Calculations of vorticity seem to be possible are possible (and interesting themselves)




# Observation of GCVE

---

- Sign of topological field fluctuations unknown – need quadratic (in induced current) effects
- CME – like-sign and opposite-sign correlations – S. Voloshin
- No antineutrons, but like-sign baryonic charge correlations possible
- Look for neutron pairs correlations!
- MPD may be well suited for neutrons!

# Estimates of statistical accuracy at NICA MPD (months of running)


- UrQMD model :  $Au + Au$  at  $\sqrt{s_{NN}} = 9$  GeV
- 2-particles  $\rightarrow$  3-particles correlations  
no necessity to fix  
the event plane
- 2 neutrons from  
mid-rapidity ( $|\eta| < 1$ )
- +1 from ZDC ( $|\eta| > 3$ )



# Other sources of quadratic effects

---

- Quadratic effect of induced currents – not necessary involve (C)P-violation
- May emerge also as C&P even quantity
- Complementary probes of two-current correlators desirable
- Natural probe – dilepton angular distributions



# Observational effects of current correlators in medium

---

- McLerran Toimela'85  $W^{\mu\nu} = \int d^4x e^{-iq \cdot x} \langle J^\mu(x) J^\nu(0) \rangle$
- Dileptons production rate

$$\begin{aligned} \frac{d(R/V)}{d^4q d^3p d^3p'} &= - \frac{1}{E_p E_{p'}} e^4 \frac{1}{(2\pi)^6} \\ &\times \delta^{(4)}(p + p' - q) L^{\mu\nu}(p, p') \\ &\times (1/q^4) W_{\mu\nu}(q) . \end{aligned}$$


- Structures –similar to DIS F1, F2  
(p → v)


# Tensor polarization of in-medium vector mesons (Bratkovskaya, Toneev, OT'95)

- Hadronic in-medium tensor – analogs of spin-averaged structure functions:  
 $p \rightarrow v$
- Only polar angle dependence
- Tests for production mechanisms

$$W^{\mu\nu} = W_1(q^2, vq) \left( g^{\mu\nu} - \frac{q^\mu q^\nu}{q^2} \right) + W_2(q^2, vq) \left( v^\mu - q^\mu \frac{vq}{q^2} \right) \left( v^\nu - q^\nu \frac{vq}{q^2} \right)$$

$$\frac{d\sigma}{d\cos\theta} \sim 1 + \frac{|v|^2}{2W_1/W_2 + 1 - (vq)^2/q^2} \cos^2\theta$$





# Effect of EM fields

---

- New structures
 
$$\begin{aligned}
 & W_1(-g^{\mu\nu} + q^\mu q^\nu / q^2) + W_2 \tilde{v}^{\mu} \tilde{v}^{\nu} \quad (\tilde{v} = v - q(vq) / q^2) \\
 & + W_3(F\tilde{F}, (F\tilde{F})^2) F^{q\mu} \tilde{v}^{\nu} + (\mu < - > \nu) + W_4 \tilde{F}^{\mu} \tilde{v}^{\nu} + (\mu < - > \nu) \\
 & + W_5 F^{q\mu} F^{q\nu} + W_6 \tilde{F}^{\mu} \tilde{F}^{\nu} + W_7 (F\tilde{F}) F^{q\mu} \tilde{F}^{\nu} + (\mu < - > \nu)
 \end{aligned}$$
- CG type relations in the real photon limit
 
$$W_2 = -W_1 q^2 / (vq)^2, W_1 \sim const$$
- Linear terms – zero real photon limit
 
$$W_3, W_4 \sim q^2 const$$
- Effect on polar and **azimuthal** asymmetries – in progress (V. Shmakova, OT)

# General hadronic tensor and dilepton angular distribution

- Angular distribution

$$d\sigma \propto 1 + \lambda \cos^2 \theta + \mu \sin 2\theta \cos \phi + \frac{\nu}{2} \sin^2 \theta \cos 2\phi + \rho \sin 2\theta \sin \phi + \sigma \sin^2 \theta \sin 2\phi$$

- Positivity of the matrix (= hadronic tensor in dilepton rest frame)

$$\begin{pmatrix} \frac{1-\lambda}{2} & \mu & \rho \\ \mu & \frac{1+\lambda+\nu}{2} & \sigma \\ \rho & \sigma & \frac{1+\lambda-\nu}{2} \end{pmatrix} \quad \begin{aligned} |\lambda| \leq 1, \quad |\nu| \leq 1 + \lambda, \quad \mu^2 &\leq \frac{(1-\lambda)(1+\lambda-\nu)}{4} \\ \rho^2 &\leq \frac{(1-\lambda)(1+\lambda+\nu)}{4}, \quad \sigma^2 \leq \frac{(1-\lambda)^2 - \nu^2}{4} \end{aligned}$$


- + cubic – det M > 0

- 1<sup>st</sup> line – Lam&Tung by SF method

# Magnetic field conductivity and asymmetries

---


- Magnetic field along z: zz-component of conductivity ( $\sim$ hadronic) tensor dominates
- Dilepton at rest:  $\lambda = -1 \rightarrow$
- Longitudinal polarization with respect to magnetic field axis
- Effects of dilepton motion – work in progress



# Other signals of rotation

---

- Hyperons (in particular,  $\Lambda$ ) polarization (self-analyzing in weak decay)
- Searched at RHIC (S. Voloshin et al.) – oriented plane (slow neutrons) - no signal observed
- No tensor polarizations as well



# Why rotation is not seen?


---

- Possible origin – distributed orbital angular momentum and local spin-orbit coupling
- Only small amount of collective OAM is coupled to polarization
- The same should affect lepton polarization
- Global (pions) momenta correlations (handedness)

# New sources of $\Lambda$ polarization coupling to rotation

---


- Bilinear effect of vorticity (HD helicity –  $\mathbf{v} \text{ curl } \mathbf{v}$ )
- generates quark axial current (Son, Surowka)
- Strange quarks - should lead to  $\Lambda$  polarization
- Proportional to chemical potential – small at RHIC – may be probed at FAIR & NICA



# Anomaly for massive quarks

---

- One way of calculation – finite limit of regulator fermion contribution (to TRIANGLE diagram) in the infinite mass limit
- The same (up to a sign) as contribution of REAL quarks
- For HEAVY quarks – cancellation!
- Anomaly – violates classical symmetry for massless quarks but restores it for heavy quarks



# Heavy quarks polarisation

---

- Non-complete cancellation of mass and anomaly terms (97)


$$\partial^\mu j_{5\mu}^c = \frac{\alpha_s}{48\pi m_c^2} \partial^\mu R_\mu, \quad \langle N(\mathbf{p}, \lambda) | j_{5\mu}^{(c)}(0) | N(\mathbf{p}, \lambda) \rangle$$

$$= \frac{\alpha_s}{12\pi m_c^2} \langle N(\mathbf{p}, \lambda) | g \sum_{f=u,d,s} \bar{\psi}_f \gamma_\nu \tilde{G}_{\mu\nu} \psi_f | N(\mathbf{p}, \lambda) \rangle$$

$$R_\mu = \partial_\mu (G_{\rho\nu}^a \tilde{G}^{\rho\nu,a}) - 4(D_\alpha G^{\nu\alpha})^a \tilde{G}_{\mu\nu}^a$$

$$= \frac{\alpha_s}{12\pi m_c^2} 2m_N^3 \delta_{\mu 3} f_S^{(2)}.$$

- Gluons correlation with nucleon spin – twist 4 operator NOT directly related to twist 2 gluons helicity BUT related by QCD EOM to singlet twist 4 correction (colour polarisability) f2 to g1
- “Anomaly mediated” polarisation of heavy quarks




# Numerics

---

- Small (intrinsic) charm polarisation

$$\bar{G}_A^c(0) = -\frac{\alpha_s}{12\pi} f_S^{(2)} \left( \frac{m_N}{m_c} \right)^2 \approx -5 \times 10^{-4}$$


- Consider STRANGE as heavy! –  
CURRENT strange mass squared is  
 $\sim 100$  times smaller – -5% -  
reasonable compatibility to the data!  
(But problem with DIS and SIDIS)



# Can $s$ REALLY be heavy?!

---


- Strange quark mass close to matching scale of heavy and light quarks – relation between quark and gluon vacuum condensates (similar cancellation of classical and quantum symmetry violation – now for trace anomaly). BUT - common belief that strange quark cannot be considered heavy,
- In nucleon (no valence “heavy” quarks) rather than in vacuum - may be considered heavy in comparison to small genuine higher twist – multiscale nucleon picture



# Comparison : Gluon Anomaly for massless and massive quarks

---

- Mass independent
- Massless (Efremov, OT '88) – naturally (but NOT uniquely) interpreted as (on-shell) gluon circular polarization
- Small gluon polarization – no anomaly?!
- Massive quarks – acquire “anomaly polarization”
- May be interpreted as a sort of correlation of quark current to chromomagnetic field
- Qualitatively similar to CME
- Very small numerically
- Small strange mass – partially compensates this smallness and leads to % effect



# Heavy unpolarized Strangeness: vector current

---

- Follows from Heisenberg-Euler effective lagrangian  
Published in **Z.Phys.98:714-732,1936.**  
e-Print: **physics/0605038**
- FFFF -> FG GG -> Describes strangeness contribution to nucleon magnetic moment and pion mean square radius
- FFFF->FFGG -> perturbative description of chiral magnetic effect for heavy (strange) quarks in Heavy Ion collisions – induced current of strange quarks

# Induced current for (heavy - with respect to magnetic field strength) strange quarks

- Effective Lagrangian

$$L = c(F\tilde{F})(G\tilde{G})/m^4 + d(FF)(GG)/m^4$$

- Current and charge density from  $c$  ( $\sim 7/45$ ) – term  $j^\mu = 2c\tilde{F}^{\mu\nu}\partial_\nu(G\tilde{G})/m^4$
- $\rho \sim \vec{H}\vec{\nabla}\theta$  (multiscale medium!)  
 $\theta \sim (G\tilde{G})/m^4 \rightarrow \int d^4x G\tilde{G}$
- Light quarks -> matching with D. Kharzeev et al' -> correlation of density of electric charge with a gradient of topological one (Lattice ?)


# Properties of perturbative charge separation

- Current carriers are obvious - strange quarks -> matching -> light quarks?
- NO obvious relation to chirality – contribution to axial current starts from pentagon (!) diagram
- No relation to topology (also pure QED effect exists)
- Effect for strange quarks is of the same order as for the light ones if topological charge is localized on the distances  $\sim 1/m_s$ , strongly (4<sup>th</sup> power!) depends on the numerical factor : Ratio of strange/light – sensitive probe of correlation length
- Universality of strange and charm quarks separation - charm separation suppressed as  $(m_s / m_c)^4 \sim 0.0001$
- Charm production is also suppressed – relative effects may be comparable at moderate energies (NICA?) – but low statistics

# Comparing CME to strangeness polarization

---


- **Strangeness polarization** – correlation of
  - (singlet) quark current
  - (chromo)magnetic field
  - (nucleon) helicity
- **Chiral Magnetic Effect** - correlation of
  - (electromagnetic) quark current
  - (electro)magnetic field
  - (Chirality flipping) Topological charge gradient



# Local symmetry violation

---

- CME – assumed to be the sign of local  $P(C)$  violation
- BUT Matrix elements of topological charge, its density and gradient are zero
- Signs of real  $C(P)$  violation – forbidden processes




# Forbidden decays in vacuum – allowed in medium

---


- C-violation by chemical potential -> (Weldon '92)  $\sigma \rightarrow e^+ e^-$
- $\rho \rightarrow \gamma\gamma$  (OT'96; Radzhabov, Volkov, Yudichev '05,06 - NJL)
- New (?) option:  $\pi \rightarrow e^+ e^-$  in magnetic field  $\frac{\Gamma_{\pi \rightarrow e^+ e^-}}{\Gamma_{\pi \rightarrow \gamma\gamma}} \sim \frac{H^2}{m_\pi^4}$
- Polarization (angular distribution in c.m. frame) of dilepton  $\sim 1 + \cos^2 \theta$  (with respect to field direction!)

# Approximation: EM part – vacuum value Two-stage forbidden decays - I



# Two-stage forbidden decays -

II




# Relating forbidden and allowed decays

- In the case of complete mass degeneracy (OT'05, unpublished):

$$\frac{\Gamma_{\sigma \rightarrow e^+ e^-}}{\Gamma_{\rho \rightarrow \gamma\gamma}} = \frac{9}{4} \frac{\Gamma_{\rho \rightarrow e^+ e^-}}{\Gamma_{\sigma \rightarrow \gamma\gamma}}$$

- Tests and corrections – in progress



# Conclusions

---

- Axial anomaly in medium is a fundamental property of QCD and may be manifested in the angular and spin asymmetries
- Chiral Vortaic Effect may be probed in the neutron asymmetries at NICA
- Bilinear current correlator may be probed in dilepton asymmetries
- CME/CVE for (heavy) strange quarks is similar to their polarization in a nucleon
- Various medium-induced decays may be related to each other