


MPD plenary meeting  
March 2017

# Femtoscscopy study for NICA / MPD

P. Batyuk (JINR), Iu. Karpenko (BITP, Kiev & INFN, Florence) ,  
R. Lednicky (JINR), L. Malinina (SINP MSU- JINR),  
K. Mikhaylov (ITEP & JINR), O. Rogachevsky (JINR),  
D. Wielanek (WUT, Warsaw)

# Introduction

**Correlation femtoscopy** : measurement of space-time characteristics  $R, c\tau \sim \text{fm}$  of particle production using particle correlations due to the effects of quantum statistics ( **QS** ) and final state interactions ( **FSI** )




- **Two particle Correlation Function (CF):**

Theory: 
$$C(q) = \frac{N_2(p_1, p_2)}{N_1(p_1) \cdot N_2(p_1)}, C(\infty) = 1$$

Experiment: 
$$C(q) = \frac{S(q)}{B(q)}, q = p_1 - p_2$$

$S(q)$  – pairs from same event  
 $B(q)$  – pairs from different event

- **Parametrization:**



**1D:**  $C(q_{inv}) = 1 + \lambda \exp(-R^2 q_{inv}^2)$  **R** Gaussian radius in Pair Rest Frame (**PRF**),  $\lambda$  correlation strength parameter

**3D:** 
$$C(q_{out}, q_{side}, q_{long}) = 1 + \lambda \exp(-R_{out}^2 q_{out}^2 - R_{side}^2 q_{side}^2 - R_{long}^2 q_{long}^2)$$


where both **R** and **q** are in Longitudinally Co-Moving Frame (**LCMS**)  
 long || beam; out || transverse pair velocity  $\mathbf{v}_T$ ; side normal to out, long

# Expected features of 1<sup>st</sup> order PT


STAR, Phys.Rev. C92 (2015) 1, 014904

- It was predicted that for 1<sup>st</sup> order phase transition  $R_{out}/R_{side} > 1$  & large  $R_{long}$  due to emission stalling during phase transition

( S. Pratt, Phys. Rev. D 33 (1986) 1314. G. Bertsch, M. Gong, M. Tohyama, Phys. Rev. C 37 (1988) 1896  
 D. H. Rischke and M. Gyulassy, Nucl. Phys. A608, 479 (1996)




- But r-t correlations in expanding source reduce the observed  $R_{out} \rightarrow R_{out}/R_{side}$
- What do the modern hydrodynamic (hybrid) models expect ?




# Expected features of 1<sup>st</sup> order PT

STAR, Phys.Rev. C92 (2015) 1, 014904

## Critical point ?



- A peak in  $R_{out}/R_{side}$  may not be related with CEP
- What do the modern hydrodynamic (hybrid) models expect ?
- VHLLE+UrQMD model was used for this study



# ArXiv 1703.09628

## Correlation femtoscopy study at NICA and STAR energies within the vHLLE+UrQMD model

P. Batyuk,<sup>1,\*</sup> Iu. Karpenko,<sup>2,3</sup> R.Lednický,<sup>1</sup> L.Malinina,<sup>1,4,5</sup> K.Mikhaylov,<sup>1,6</sup> O. Rogachevsky,<sup>1</sup> and D.Wielanek<sup>7</sup>

<sup>1</sup>*Veksler and Baldin Laboratory of High Energy Physics, JINR Dubna, 141980 Dubna, Russia*

<sup>2</sup>*Bogolyubov Institute for Theoretical Physics, 03680 Kiev, Ukraine*

<sup>3</sup>*INFN - Sezione di Firenze, I-50019 Sesto Fiorentino (Firenze), Italy*

<sup>4</sup>*M. V. Lomonosov Moscow State University, Moscow, Russia*

<sup>5</sup>*D. V. Skobeltsyn Institute of Nuclear Physics, Moscow, Russia*

<sup>6</sup>*Institute of Theoretical and Experimental Physics (ITEP), Moscow, Russia*


<sup>7</sup>*Warsaw University of Technology, Faculty of Physics, Warsaw 00662, Poland*

Correlation femtoscopy allows one to measure the space-time characteristics of particle production in relativistic heavy-ion collisions due to the effects of quantum statistics (QS) and final state interactions (FSI). The main features of the femtoscopy measurements at top RHIC and LHC energies are considered as a manifestation of strong collective flow and are well interpreted within hydrodynamic models employing equation of state (EoS) with a crossover type transition between Quark-Gluon Plasma (QGP) and hadron gas phases. The femtoscopy at lower energies was intensively studied at AGS and SPS accelerators and is being studied now in the Beam Energy Scan program (BES) at the BNL Relativistic Heavy Ion Collider in the context of exploration of the QCD phase diagram. In this article we present femtoscopic observables calculated for Au-Au collisions at  $\sqrt{s_{NN}} = 7.7 - 62.4$  GeV in a viscous hydro + cascade model vHLLE+UrQMD and their dependence on the EoS of thermalized matter.

PACS numbers: 25.75.-q, 25.75.Gz

Keywords: relativistic heavy-ion collisions, hydrodynamics, collective phenomena, Monte Carlo simulations, vHLLE, UrQMD

# vHLLE+UrQMD model




Iu. Karpenko, P. Huovinen, H.Petersen, M. Bleicher, Phys.Rev. C 91, 064901 (2015), arXiv:1502.01978,1509.3751, talk QM2015  
 vHLLE code: free and open source, <https://github.com/yukarpenko/vhllle>, Comput. Phys. Commun. 185 (2014), 3016

Model tuned by matching with the experimental data of SPS and BES RHIC.


Chiral EoS → crossover PT (XPT)  
 J. Steinheimer, et al, J. Phys. G 38, 035001 (2011)

HadronGas + Bag Model → 1<sup>st</sup> order PT (1PT)  
 P.F. Kolb, et al, Phys.Rev. C 62, 054909 (2000)

Pion emission times at the particlization surface



and at the last interaction points






# vHLE+UrQMD model

Mean CMS times of pion emission at  
particlization and last interaction

$\sqrt{s_{NN}}$ [GeV]	EoS	particlization surface		last interactions	
		$\bar{t}$ [fm/c]	RMS [fm/c]	$\bar{t}$ [fm/c]	RMS [fm/c]
7.7	1PT	7.24	2.84	13.15	6.56
	XPT	6.16	2.01	11.61	6.26
11.5	1PT	7.33	2.31	13.09	6.92
	XPT	6.36	1.91	11.57	6.41
19.6	1PT	6.88	2.16	13.18	7.56
	XPT	6.41	2.15	11.93	6.93
27	1PT	6.85	2.37	13.38	8.07
	XPT	6.40	2.39	12.62	7.57
39	1PT	7.17	2.75	13.98	8.30
	XPT	6.64	2.58	13.05	7.85
62.4	1PT	7.00	2.82	14.11	8.50
	XPT	6.60	2.63	12.72	7.81

CMS times of pion emission at last interaction






Emission times for 1<sup>st</sup> order phase transition  
are larger than for crossover.

- Weak dependence of the average pion creation time on the collision energy.  
Maximal difference :  $\sim 1.5$  fm. Interplay of longer pre- thermal and shorter hydrodynamic stage at lower collision energies
- On the other hand, the duration of hydro stage gets shorter as collision energy decreases because of lower initial energy density at the hydro starting time.
- The cascade smears the relative difference between the 1PT and XPT scenarios

We are studying the possibilities to extract this difference experimentally at NICA/MPD using femtoscopy technique.


# Correlation functions with vHLE+UrQMD

- The difference between pion CF for 1<sup>st</sup> order PT and crossover < 5%
- For kaons it is expected to be larger ~ 10%
- It is necessary to study different particle types. Importance of PID





# 3D Pion radii versus $m_T$ with vHLLE+UrQMD model




Green triangles - 1PT EoS, Red triangles - XPT EoS, Open black squares STAR data BES


- $R_{out}$  (XPT) at high energies and  $R_{out}$  (1PT) at all energies are slightly overestimated -> an indication of the need to reduce the emission time in the model
- $R(1PT) > R(XPT)$  by  $\sim 1$  fm for “out” and “long” radii

# $R_{out}/R_{side}$ with vHLLE + UrQMD model

- $R_{out}/R_{side}$  and  $R_{out} - R_{side}$  as a function of  $s_{NN}$  were studied at fixed  $m_T$  by the STAR Collaboration. A wide maximum near  $s_{NN} \sim 20$  GeV/c in both excitation functions was observed.






- $R_{out}/R_{side}$  (XPT) agrees with almost all STAR data points within rather large systematic errors, while  $R_{out}/R_{side}$  (1PT) overshoots the data.



There is an indication in our study that optimal description of the femtoscopic radii requires about 1 fm shorter duration of pion emission with the present setup of the model, at all collision energies. It is an open question whether a new set of parameters can be found which accommodates the the femtoscopic radii.

# Ratio $R_{osl}^{1PT} / R_{osl}^{XPT}$ versus $\sqrt{s_{NN}}$



- $R_{side}$  radii in the 1PT EoS and XPT EoS scenarios practically coincide;
- $R_{out}$  ( $R_{long}$ ) for 1PT EoS > XPT EoS, strong dependence on  $k_T$  interval
- The difference comes from weaker transverse flow developed in the fluid phase with 1PT EoS as compared to XPT EoS & longer lifetime of the fluid phase in 1PT EoS

# Source functions

The new Source Function technique was used.

SF for 1<sup>st</sup> order is wider than the one for crossover.

Main advantage of this technique is the possibility to use the Source Functions itself without any hypothesis about its shape.


$$C(\mathbf{k}^*, \mathbf{P}) = \int d^3\mathbf{r}^* S^\alpha(\mathbf{r}^*, \mathbf{P}) \left| \psi_{-\mathbf{k}^*}^{S, \alpha' \alpha}(\mathbf{r}^*) \right|^2,$$

Different functions were tested to describe the shape of SF projections: single Gaussian

$$S(\vec{r}^*) \sim \exp\left(-\frac{r_{out}^{*2}}{4R_{out}^{*2}} - \frac{r_{side}^{*2}}{4R_{side}^{*2}} - \frac{r_{long}^{*2}}{4R_{long}^{*2}}\right),$$

$$S^H(r_x, r_y, r_z) = \lambda \exp\left[-f_s\left(\frac{x^2}{4r_{xs}^2} + \frac{y^2}{4r_{ys}^2} + \frac{z^2}{4r_{zs}^2}\right) - f_l\left(\frac{x^2}{4r_{xl}^2} + \frac{y^2}{4r_{yl}^2} + \frac{z^2}{4r_{zl}^2}\right)\right],$$

$$f_s = 1/[1 + (r/r_0)^2], \quad f_l = 1 - f_s.$$




Gaussian + Gaussian;  
Gaussian + Lorencian  
Gaussian + Exp

Hump-function


- The best description was obtained with Gaussian+Gaussian and Hump-function.
- Gaussian+Gaussian - simple interpretation (core-resonances) & more stable fitting procedure

# Source Function with vHLLE + UrQMD model



- Two-Gaussian fit describes reasonably SF till  $\sim 60$  fm «out» and  $\sim 25$  fm «side» and «long» directions.
- One-Gaussian fit gives large  $\chi^2 / \text{NdF}$ , but the values of radii are equal to the ones of two Gaussian radii averaged according with relative contributions of small and large radii; **That is why it reflects reasonably the main features of 2-Gaussian fit at small  $r^*$ .**

# Pion Source Function with $\nu$ HLE + UrQMD



- “out” :  $R_{out,2}$  (1PT)  $>$   $R_{out,2}$  (XPT); for the calculations with the first order phase transition and for the one with crossover phase transition decreases with increasing  $\sqrt{s_{NN}}$  ;

The relative contributions of small and large radii

$\lambda_1 \sim 0.65$  and  $\lambda_2 \sim 0.35$  do not depend on  $\sqrt{s_{NN}}$  and on the type of EoS.

- The radii  $R_{side,2}$  and corresponding  $\lambda_{1,2}$  do not depend on  $\sqrt{s_{NN}}$  .
- “long”: radii almost coincide for both types of EoS, The relative contribution of the large radii,  $\lambda_2$  increases with  $\sqrt{s_{NN}}$  , while  $\lambda_1$  decreases.  $\lambda_2$  (1PT)  $>$   $\lambda_2$  (XPT)




# Conclusions

- Possibility to distinguish between hybrid model source functions with 1<sup>st</sup> order phase transition and crossover was studied using vHLE+UrQMD model
  - Hydro phase lasts longer with 1<sup>st</sup> order PT.
  - Hadronic cascade diminishes the difference between 1PT and XPT source functions, though there is still a possibility to distinguish them using the femtoscopy technique.
  - vHLE+UrQMD model with XPT describes RHIC femtoscopy radii at  $\sqrt{s_{NN}} = 7.7-62.4$  GeV
  - There is an indication that optimal description of the femtoscopy radii requires about 1 fm shorter pion emission time with the present setup of the model, at all collision energies. - new tune of vHLE+UrQMD model is needed.
- It'll be very interesting to try to use 3 phase hydro model (THESEUS) at low energies
- $R_{out}(1PT) > R_{out}(XPT)$  &  $R_{long}(1PT) > R_{long}(XPT)$
  - -----
  - Source functions technique allows to get an additional information about differences between 1PT / XPT; Best parametrizations of SF : Gauss+Gauss and Hump
  - The standard one-Gaussian parametrization of the 3D CF reflects correctly the behaviour of the SF at small  $r^*$  and is sufficiently sensitive to EoS.
  - It is very promising to make 3D CF analysis using heavier particles: K,p because of more Gaussian shape of SF and less influence of resonances


# MPD detector has the same advantages as ALICE to study femtoscopy:

- It can be promising to make 3D CF analysis using heavier particles: K, p because of more Gaussian shape of SF and less influence of resonances
- Different particle pairs:  $\pi K$ ,  $K+K^-$ ,  $\pi p$ ,  $\pi \Lambda$ ,  $\Lambda \Lambda$  .. can be studied -- different influence of cascade phase, emission assymetries..
- Az-sensitive femtoscopy is particularly sensitive to the evolution time (in addition to  $R_{long}$ ) and to the expansion velocity.

ALICE	MPD
<ul style="list-style-type: none"> <li>• Low momentum cut-off (<math>p_T &gt; 100 \text{ MeV}/c</math>)</li> <li>• Small material budget</li> <li>• Excellent particle identification (PID) by: specific energy loss (<math>dE/dx</math>) &amp; TOF</li> <li>• Good primary and secondary vertex resolution</li> </ul>	<ul style="list-style-type: none"> <li>✓</li> <li>✓</li> <li>✓</li> <li>✓</li> </ul>



In femtoscopy context, the remarkable feature of collider experiments, compared with fixed target ones, is an excellent particle identification (not so important for correlations of identical pions, while quite important for other particle pairs) and a good two-track resolution.



*Back Up*


# Imaging

- PHENIX and STAR collaborations apply a new “imaging technique” to extract the  $S(r^*)$ -source function, which represents time-integrated distribution of particle emission points separation  $r^*$  in the pair rest frame (PRF).

$$C(\mathbf{q}) - 1 \equiv R(\mathbf{q}) = \int (|\phi(\mathbf{q}, \mathbf{r})|^2 - 1) S(\mathbf{r}) d\mathbf{r},$$

- The method is suitable for extracting the  $S(r)$  directly from the data without any hypothesis about source shape; it seems to be very useful for comparison of the experimental data with the models with 1PT or Crossover EoS
- The good knowledge of all factors influencing the shape of correlation function is needed

STAR, Phys.Rev. C88 (2013) 3, 034906




# Present activities of SINP MSU group

within the agreement between JINR and SINP MSU : 01.06.16-30.04.17

---

- **Development of tracking algorithm for ITS MPD**  
(talk of Stanislav Shushkevich)
- **Femtoscopia study for NICA .**
  - MC study of femto observables
  - PID & tracking studies (talk of Gulnara Eyyubova)
  - Development of MPD Femto software  
(Malinina Ludmila, Konstantin Mikhailov, Gulnara Eyyubova,  
together with JINR group of Oleg Rogachevsky)
- **Alignment for BM@N** (talk of Anatoly Solomin)


# UrQMD 3.4 model



H. Petersen, J. Steinheimer, G. Burau, M. Bleicher and H. Stöcker, Phys. Rev. C 78 (2008) 044901.  
UrQMD-3.4 code was taken from <http://urqmd.org/>  
Many thanks to Hannah Petersen for the advises concerning parameters of simulations!

- Initial collisions and string fragmentations from the microscopic UrQMD model.
- (3+1)-dimensional ideal hydrodynamic evolution.
- hadronic cascade.

Chiral EoS - Crossover  
Bag model EoS - 1st order  
Hadron gas EoS




Both models describe the bulk data quite well, but the vHLL+UrQMD seems to have a better description whereas UrQMD 3.4 tends to overestimate the femtoscopic radii.




# Kaon Source Function with vHLE + UrQMD

Kaons 7.7 GeV/c Gauss+Gauss fit of 3D SF projections kT (0.2-0.8) GeV/c



- “out” : for the same  $k_T$  range the difference 1PT/XPT is smaller for heavier particles than for pions due to weaker influence of flow effects
- For kaons: more clear signal, less influence of resonances, --> more Gaussian shape

# Femtoscscopy



**Correlation femtoscopy** : measurement of space-time characteristics  $R, c\tau \sim \text{fm}$  of particle production using particle correlations due to the effects of **QS** and **FSI**

G. Goldhaber, S. Goldhaber, W-Y Lee, A. Pais (Phys.Rev. 120 (1960) 300): first showed the BE correlation of identical pions in  $pp^-$  collisions

G.I. Kopylov and M.I. Podgoretsky (1971-1975) (review: Phys.Part.Nucl. 20, iss. 3 (1989) 629, in Russian): elaborated [basics of correlation femtoscopy](#)

V.G. Grishin, G.I. Kopylov, and M.I. Podgoretsky showed [analogy](#) (Sov.J.Nucl.Phys. 13 (1971) 638) and [difference](#) (G.I. Kopylov and M.I. Podgoretsky, Sov.J.Nucl.Phys. 15 (1972) 219) between [femtoscscopy in particle physics](#) and [HBT effect in astronomy](#) (R. Hanbury-Brown and R.Q. Twiss, Phil.Mag. 45 (1954) 633): [HBT effect](#) is the change of intensity of the signal received from the particle emission source