

Black stars induced by matter on a brane: exact solutions

Maxim Kurkov

in collaboration with Prof. A.A. Andrianov

based on: arXiv:1008.2705

V.A.Fock Department of Theoretical Physics
Saint-Petersburg State University

International Workshop Bogoliubov Readings
Russia, Dubna, 2010

Introduction

- What is the problem?
 - **Black hole** creation may be a consequence of strong gravity at short distances attainable in high energy experiments if our space is realized on a hypersurface – three-brane in a multidimensional space-time.
 - Correct (or better exact) description of black hole geometry when the **matter universe is strictly situated on the three-dimensional brane but gravity propagates into extra space dimensions** is needed.
 - Appearance of **delta-like singularities in matter distribution** hidden under horizon for static locally stable black holes is a problem: in fact the **matter (quarks and gluons) must be smoothly distributed**.
 - Therefore one expects that rather **black stars** are created with matter both inside and outside an event horizon in a finite brane-surface volume.

Techniques

- Stress-energy tensor structure.

- The Einstein equations in the bulk read,

$${}^{(5)}G_{AB} = \kappa_5 T_{AB}, \quad T_{AB} = \delta_A^\mu \delta_B^\nu \tau_{\mu\nu} \delta(z)$$

with $\kappa_5 = 1/M_*^3$ and M_* is a Planck scale in five dimensions.

- In order to define $\tau_{\mu\nu}$ let us introduce extrinsic curvature tensor $K_{\mu\nu}$.

$$K_{\mu\nu} = -\frac{1}{2} \frac{\partial g_{\mu\nu}}{\partial z} \text{ valid in the Gaussian normal coordinates (i.e. } g_{zz} = -1, g_{\mu z} = 0 \text{) only!}$$

- $\tau_{\mu\nu}$ is defined by the Israel-Lanczos junction conditions,

$$[g_{\mu\nu} K - K_{\mu\nu}]_{-0}^{+0} = \kappa_5 \tau_{\mu\nu}.$$

- $K_{\mu\nu}^{+0}$, and $K_{\mu\nu-0}$ are the extrinsic curvature tensors of hypersurfaces $z = +0$ and $z = -0$ correspondingly.

Techniques

- General construction.
 - To build a brane we search for a metric $g_{AB}(x, y)$ which is a bulk vacuum solution of the Einstein equations with event horizon.
 - Suppose that:
 - a) the induced metric $g_{\mu\nu}(x, y)$ is asymptotically flat for any hypersurface $y = \text{const}$ and inherits the horizon;
 - b) in the chosen coordinate systems $g_{5B}(x, y) = 0$ and the remaining metric components provide orbifold geometry $g_{AB}(x, y) = g_{AB}(x, -y)$;
 - c) Coordinate y is spacelike i.e. $g_{yy} \equiv g_{55} < 0$.
 - In order to generate a brane filled by matter we proceed the following transformation,

$$g_{AB}(x, y) \implies g_{AB}(x, |z| + a).$$

- Brane: $z = 0$.

Construction of the solution.

- Preparing of the suitable coordinate system.
 - We start from the metric describing a five-dimensional static neutral black hole in Schwarzschild coordinates $\{t, r, \theta_1, \theta_2, \theta\}$,

$$g_{AB} = \text{diag} \left[U(r), -\frac{1}{U(r)}, -r^2 \cos^2 \theta, -r^2 \cos^2 \theta \cos^2 \theta_1, -r^2 \right],$$

where $U(r) = 1 - \frac{M}{r^2}$, M is related to the Schwarzschild-Tangherlini radius $M \equiv r_{Sch-T}^2$.

- Let's define the Gaussian normal coordinates in respect to hypersurface with space-like normal vector $\theta = 0$.
- The vector orthonormal to this hypersurface $n^A = [0, 0, 0, 0, 1/r]$.
- The required change of coordinates acts on two variables $r = r(\rho, y)$,
 $\theta = \theta(\rho, y)$.

Construction of the solution.

- Our coordinate transformation has the following form.

-

$$|y| = \int_{\rho}^r \frac{\text{sign}((r - \rho)) x^2}{\sqrt{(x^2 - M)(x^2 - \rho^2)}} dx, \quad \theta = \int_{\rho}^r \frac{\text{sign}((r - \rho)y)}{\sqrt{(x^2 - M)(x^2 - \rho^2)}} dx.$$

- We have: inside the horizon $r < \rho < \sqrt{M}$ and outside the horizon $\sqrt{M} < \rho < r$.

- The metric in new coordinates $\{t, \rho, \theta_1, \theta_2, y\}$, reads,

$$g_{AB}(x, y) = \text{diag} \left[U(r), -\frac{r^2 r_{\rho}^2}{\rho^2 U(r)}, -r^2 \cos^2 \theta, -r^2 \cos^2 \theta \cos^2 \theta_1, -1 \right],$$

where $r = r(\rho, y)$, $\theta = \theta(\rho, y)$.

- The final answer for black star metric and $\tau_{\mu\nu}$ has the following form:

$$g_{AB}^{\text{final}}(x, z) = g_{AB}(x, y)|_{y=|z|+a}, \quad \kappa_5 \tau_{\mu\nu}(x, a) = \left(\frac{\partial g_{\mu\nu}}{\partial y} - g_{\mu\nu} \frac{g^{\lambda\delta} \partial g_{\lambda\delta}}{\partial y} \right) \Big|_{y=a}$$

Construction of the solution.




Figure 1: Pairs of hypersurfaces symmetric in respect to the horizontal axis to be glued into a brane are shown by red curves. The circle of horizon in dim = 5 is depicted by green line. $g_{AB}(x, y) \implies g_{AB}(x, |z| + a)$, $a = 0.69868\sqrt{M} \div \pi\sqrt{M}/2$

Construction of the solution.

- Some technical remarks.
 - Note, that situation on the horizon is O.K. All quantities that must be continuous are continuous. for example for scalar curvature on the brane ${}^{(4)}R$ we have the following limit:

$${}^{(4)}R(a) = -2 \frac{B + 1 - \cos^2 a - 4 |\sin a| \sqrt{1+B} \sqrt{B} \cos a}{(1+B) \cos^2 a},$$

$$B(a) \equiv \lim_{\rho \rightarrow \sqrt{M}} \frac{r(\rho, a) - \rho}{\rho - \sqrt{M}} = \frac{1}{2} \left(\cosh \left(\frac{2a}{\sqrt{M}} \right) - 1 \right).$$

- In this construction space-time is asymptotically flat and the following asymptotic takes place:

$${}^{(4)}R = \frac{4M^2 a^2}{\rho^8} \left(1 + O \left(\frac{a^2}{\rho^2} \right) \right).$$

Matter distribution.

- Effective 4-D stress-energy tensor $S_{\mu\nu}$.

- projection of Einstein equations onto the brane: SMS equations

$${}^{(4)}G_{\mu\nu} \equiv G_{\mu\nu} = \kappa_5^2 \Sigma_{\mu\nu} - E_{\mu\nu} \equiv \kappa_4 S_{\mu\nu}, \quad \kappa_4 \equiv \frac{1}{M_{Pl}^2},$$

- where

$$\Sigma_{\mu\nu} = \frac{1}{24} \left(-2\tau\tau_{\mu\nu} + 6\tau_{\mu}^{\sigma}\tau_{\sigma\nu} + g_{\mu\nu}(-3\tau^{\sigma\rho}\tau_{\sigma\rho} + \tau^2) \right),$$

- and

$$E_{\mu\nu} = {}^{(5)}C_{BCD}^A n_A n^C q_{\mu}^B q_{\nu}^D.$$

- Compare with 5-D Einstein equations:

$${}^{(5)}G_{AB} = \kappa_5 \delta_A^{\mu} \delta_B^{\nu} \tau_{\mu\nu} \delta(z).$$

Matter distribution.

- Here and below we use new radial coordinate $R(\rho, a) \equiv r(\rho, a) \cos \theta(\rho, a)$ on the brane.
- The total mass in 4+1 dimension is given by

$$\mathcal{M} = \frac{3}{16\pi\kappa_5} \int_{t=const} d^{(4)}V {}^{(5)}R_{AB}\xi^A m^B \equiv \int_0^\infty dR f_5(R).$$

- \mathcal{M} does not depend on the value of parameter a !
- The exact calculations show that the 3-dim Komar integral, ${}^{(4)}\mathcal{M}_{eff} = 0$.
- compare with $g_{00} - 1 = O(1/R^2)$ but not $O(1/R) \Leftarrow$ infinite size of an extra dimension.

Matter distribution.




Figure 2: The matter-density radial distributions $f_5(R, a)$ on the brane with $M = 1$ are presented by a magenta colored line. The corresponding horizons are indicated by green lines. $a_1 > a_2 > a_3 > a_4 > a_5$

Matter distribution.




Figure 3: The matter-density radial distribution $f_5(R)$ on the brane with $a = 1.1$, $M = 1$ is presented for $\kappa_5 = 1$ by a magenta colored line. The effective matter-density $f_4(R)$ is shown by blue line for the value $\kappa_4 = 50$ to compare with $f_5(R)$. The horizon is indicated by green line.

Conclusions

- Results.
 - We have shown that by cut-and-paste method in special Gaussian normal coordinates one can build the exact geometry of multidimensional black star with *horizon*, generated by a *smooth* matter distribution in our universe.
 - In our approach, for a given total mass, the profiles of available configurations for matter distribution are governed by the parameter a which is presumably related to the collision kinematics when a black object ("black hole") is created by partons on colliders.
- Generalizations.
 - charged and rotated black stars as well as black rings.
compact extra dimensions and warped geometries.

Thanks for attention!!!!